## A Data flow of Wassterstein learning for point process

Figure 4 illustrates the data flow for WGANTPP.



 $\mathbf{P}$ Figure 4: The input and output sequences are  $\zeta = \{z_1, \ldots, z_n\}$  and  $\rho = \{t_1, \ldots, t_n\}$  for generator  $g_{\theta}(\zeta) = \rho$ , where  $\zeta \sim Poisson(\lambda_z)$  process and  $\lambda_z$  is a prior parameter estimated from real data. Discriminator computes the Wassterstein distance between the two distributions of sequences  $\rho = \{t_1, t_2, \ldots\}$  and  $\xi = \{\tau_1, \tau_2, \ldots\}$ 

.

## **B** Proof that  $\|\cdot\|_{\star}$  is a norm

It is obvious that  $\|\cdot\|_{\star}$  is nonnegative and symmetric. If  $\|\xi - \rho\|_{\star} = 0$ , then  $m = n$  and there is a assignment  $\sigma$  such that  $x_i = y_{\sigma(i)}$  for all  $i = 1, \ldots, n$ .

Now we prove that  $\|\cdot\|_{*}$  has triangle inequality. WLOG, assume that  $\xi = \{x_1, \ldots, x_n\}, \rho =$  $\{y_1, \ldots, y_k\}$  and  $\zeta = \{z_1, \ldots, z_m\}$  where  $n \leq k \leq m$ . Define the permutation  $\hat{\sigma}$  on  $\{1, \ldots, k\}$  by

$$
\hat{\sigma} := \arg \min_{\sigma} \sum_{i=1}^{n} \|x_i - y_{\sigma(i)}\| + \sum_{i=n+1}^{k} \|s - y_{\sigma(i)}\| \tag{11}
$$

Then we know that

$$
\|\xi - \rho\|_{\star} = \sum_{i=1}^{n} \|x_i - y_{\hat{\sigma}(i)}\| + \sum_{i=n+1}^{k} \|s - y_{\hat{\sigma}(i)}\|
$$
 (12)

Therefore, we have that

$$
\|\xi - \zeta\|_{\star} = \min_{\sigma} \sum_{i=1}^{n} \|x_i - z_{\sigma(i)}\| + \sum_{i=n+1}^{m} \|s - z_{\sigma(i)}\|
$$
  
\n
$$
\leq \min_{\sigma} \sum_{i=1}^{n} (||x_i - y_{\hat{\sigma}(i)}|| + ||y_{\hat{\sigma}(i)} - z_{\sigma(i)}||) + \sum_{i=n+1}^{k} (||s - y_{\hat{\sigma}(i)}|| + ||y_{\hat{\sigma}(t)} - z_{\sigma(i)}||)
$$
  
\n
$$
+ \sum_{i=k+1}^{m} ||s - z_{\sigma(i)}||
$$
  
\n
$$
= ||\xi - \rho||_{\star} + \min_{\sigma} \sum_{i=1}^{k} ||y_{\hat{\sigma}(i)} - z_{\sigma(i)}|| + \sum_{i=1}^{m} ||s - z_{\sigma(i)}||
$$
\n(13)

$$
= ||\xi - \rho||_{\star} + \min_{\sigma} \sum_{i=1} ||y_{\hat{\sigma}(i)} - z_{\sigma(i)}|| + \sum_{i=k+1} ||s - z_{\sigma(i)}||
$$
  

$$
= ||\xi - \rho||_{\star} + \min_{\sigma} \sum_{i=1}^{k} ||y_i - z_{\sigma(\hat{\sigma}^{-1}(i))}|| + \sum_{i=k+1}^{m} ||s - z_{\sigma(i)}||
$$
  

$$
= ||\xi - \rho||_{\star} + ||\rho - \zeta||_{\star}
$$

where the last equality is due to the fact that the minimization is taken over all permutations  $\sigma$  of  $\{1, \ldots, m\}$ , and  $\hat{\sigma}$  is a fixed permutation of  $\{1, \ldots, k\}$  where  $k \leq m$ . This completes the proof.

## C Proposed  $\|\cdot\|_{\star}$  Distance on the Real Line

In this section, we prove that finding the distance between sequences  $\xi$  and  $\rho$ ,

$$
\|\xi - \rho\|_{\star} = \min_{\sigma} \sum_{i=1}^{n} \|x_i - y_{\sigma(i)}\|_{\circ} + \sum_{i=n+1}^{m} \|s - y_{\sigma(i)}\|,
$$
\n(14)

in the case of temporal point process in [0, T], *i.e.*,  $\xi = \{t_1 < t_2 < \ldots < t_n\}$  and  $\rho =$  $\{\tau_1 < \tau_2 < \ldots < \tau_m\}$ , reduces to

$$
\|\xi - \rho\|_{\star} = \sum_{i=1}^{n} |t_i - \tau_i| + \sum_{i=n+1}^{m} (T - y_i),\tag{15}
$$

Here, without loss of generality  $n \leq m$  is assumed. The choice of  $s = T$  is basically padding the shorter sequences with  $T$ . Given, the sequences have the same length now, we claim that the identity permutation *i.e.*,  $\sigma(i) = i$  is the minimizer in (14). We proceed by a proof by contradiction. Assume that the minimizer is NOT the identity permutation. Then, find the first i such that  $\sigma(i) \neq i$ . Then,  $\Sigma(i) = j$  where  $j > i$ . Therefore, there should be a  $k > i$  such that  $\sigma(k) = i$ . Then, if you change the permutation according to  $\sigma(i) = i$  and  $\sigma(k) = j$  the cost will change by

$$
\Delta = \underbrace{(|t_i - \tau_j| + |t_k - \tau_i|)}_{\text{for the old permutation}} - \underbrace{(|t_i - \tau_i| + |t_k - \tau_j|)}_{\text{for the new permutation}} \tag{16}
$$

Given  $i < j$  and  $i < k$ , it is easy to see that  $\Delta > 0$ . This means that we've found a better permutation which contradicts our assumption. Therefore, the optimal permutation will match the event points in an increasing order one by one.

## D Equivalence of the  $\|\cdot\|_{\star}$  Distance and Difference in Count Measures

The count measure of a temporal point process is a special case of the one defined for point processes in general space in Section 2.1. For a Borel subset  $B \subset S = [0, T)$  we have  $N(B) = \int_{t \in B} \xi(t) dt$ . With a little abuse of notation we write  $N(t) := N([0, t)) = \int_0^t \xi(t) dt$ . Figure 1 is a good guidance through this paragraph. Starting from time 0 the first gap in count measure starts from  $min(t_1, \tau_1)$  and ends in  $\max(t_1, \tau_1)$ . Therefore, there is difference equal to  $s_1 = \max(t_1, \tau_1) - \min(t_1, \tau_1) = |t_1 - \tau_1|$ in the count measure. Similarly, the second block of difference has volume of  $s_2 = |t_2 - \tau_2|$ , and so on. Finally, for  $m > n$  the  $(n + i)$ -th block make a difference of  $s_{n+i} = T - \tau_{n+i}$ . Therefore, the area (L<sub>1</sub> distance) between the two sequences is a equal to  $S = \sum_{i=1}^{m} s_i$ . On the other hand by looking (15) we observe that  $\|\xi - \rho\|_{\star} = \sum_{i=1}^{m} s_i$ . Therefore, by choice of  $s = T$  as an anchor point, the distance we have is exactly the area between the two count measures.