
Supplementary Material for “Faster Online Learning of Optimal Threshold for Consistent F-measure Optimization”

Mingrui Liu^{*†}, Xiaoxuan Zhang^{*†}, Xun Zhou[‡], Tianbao Yang[†]

[†]Department of Computer Science, The University of Iowa, Iowa City, IA 52242, USA

[‡]Department of Management Sciences, The University of Iowa, Iowa City, IA 52242, USA
mingrui-liu, xiaoxuan-zhang, xun-zhou, tianbao-yang@uiowa.edu

1 Extension to Other Metrics

In this section, we consider the extension of the proposed method to other metrics, in particular Jaccard similarity coefficient, and F_β measure.

Let us first consider Jaccard similarity coefficient (JAC) [2]:

$$\text{JAC}(f) = \frac{\int_{\mathcal{X}} \eta(\mathbf{x})f(\mathbf{x})d\mu(\mathbf{x})}{\pi + \int_{\mathcal{X}} f(\mathbf{x})d\mu(\mathbf{x}) - \int_{\mathcal{X}} \eta(\mathbf{x})f(\mathbf{x})d\mu(\mathbf{x})}.$$

Then we have

$$\frac{1}{\text{JAC}(f)} = \frac{\pi + \int_{\mathcal{X}} f(\mathbf{x})d\mu(\mathbf{x})}{\int_{\mathcal{X}} \eta(\mathbf{x})f(\mathbf{x})d\mu(\mathbf{x})} - 1 = \frac{2}{F(f)} - 1.$$

Therefore

$$\text{JAC}(f) = \frac{F(f)}{2 - F(f)}.$$

If $F(f)$ is maximized so is $\text{JAC}(f)$. According to [2], the optimal threshold $\theta_{\text{JAC},*}$ for maximizing $\text{JCA}(\eta_\theta(\mathbf{x}))$ is given by $\theta_{\text{JAC},*} = \frac{\text{JAC}_*}{1 + \text{JAC}_*} = F_*/2 = \theta_{F,*}$, where $\theta_{F,*}$ is the optimal threshold for F-measure maximization. Given an estimate of θ_F for F-measure optimization, we can set the threshold for JAC maximization as $\theta_{\text{JAC}} = \theta_F$, and when $\theta_F \rightarrow \theta_{F,*}$, we have $\theta_{\text{JAC}} \rightarrow \theta_{\text{JAC},*}$. As a result, the proposed algorithm FOFO is still applicable.

Next, let us consider F_β -measure:

$$F_\beta(f) = \frac{(1 + \beta^2) \int_{\mathcal{X}} \eta(\mathbf{x})f(\mathbf{x})d\mu(\mathbf{x})}{\beta^2\pi + \int_{\mathcal{X}} f(\mathbf{x})d\mu(\mathbf{x})}.$$

Following the same analysis as in [3][Lemma 13, Lemma 14], we can have that $F_\beta(\eta_\theta)$ is maximized at a point $\theta_{\beta,*}$ that is the root of the following equation:

$$\pi\beta^2\theta - \mathbb{E}_{\mathbf{x}}[(\eta(\mathbf{x}) - \theta)_+] = 0,$$

which is the optimal solution of the following strongly convex function

$$Q(\theta) \triangleq \frac{1}{2}\mathbb{E}_{\mathbf{x}}[(\eta(\mathbf{x}) - \theta)_+]^2 + \frac{1}{2}\pi\beta^2\theta^2.$$

*equal contribution

For the optimal threshold $\theta_{\beta,*}$ and optimal $F_{\beta,*}$, we have $\theta_{\beta,*} = \frac{F_{\beta,*}}{1+\beta^2}$. Hence, we can search for $\theta_{\beta,*}$ by solving the following problem:

$$\min_{\theta \in [0, 1/(1+\beta^2)]} Q(\theta) \triangleq \frac{1}{2} \mathbb{E}_{\mathbf{x}} [(\eta(\mathbf{x}) - \theta)_+]^2] + \frac{1}{2} \pi \beta \theta^2.$$

We can modify FOFO a little to account for this change.

2 Missing Proofs

2.1 \mathbf{w}_* minimizes the expected logistic loss

Under the assumption that

$$\eta(\mathbf{x}) = \Pr(y = 1 | \mathbf{x}, \mathbf{w}_*) = \frac{1}{1 + \exp(-\mathbf{w}_*^\top \phi(\mathbf{x}))},$$

we prove that \mathbf{w}_* is the minimizer of the following problem:

$$\min_{\mathbf{w} \in \mathbb{R}^d} L(\mathbf{w}) \triangleq \mathbb{E}_{\mathbf{x}, y} \log(1 + \exp(-(2y - 1)\mathbf{w}^\top \phi(\mathbf{x}))). \quad (7)$$

Using variable change $\tilde{y} = 2y - 1$, $\Pr(\tilde{y} | \mathbf{x}, \mathbf{w}_*) = \frac{1}{1 + \exp(-\tilde{y}\mathbf{w}_*^\top \phi(\mathbf{x}))}$, and $L(\mathbf{w}) = \mathbb{E}_{\mathbf{x}, \tilde{y}} \log(1 + \exp(-\tilde{y}\mathbf{w}^\top \phi(\mathbf{x})))$. Then,

$$\begin{aligned} L(\mathbf{w}) &= \mathbb{E}_{\mathbf{x}, \tilde{y}} \log(1 + \exp(-\tilde{y}\mathbf{w}^\top \phi(\mathbf{x}))) = - \int_{\mathbf{x}} \mathbb{E}_{\tilde{y} | \mathbf{x}} [\log \Pr(\tilde{y} | \mathbf{x}, \mathbf{w})] d\mu(\mathbf{x}) \\ &= \int_{\mathbf{x}} \left[- \sum_{\tilde{y}} \Pr(\tilde{y} | \mathbf{x}, \mathbf{w}_*) \log \Pr(\tilde{y} | \mathbf{x}, \mathbf{w}) \right] d\mu(\mathbf{x}) \end{aligned}$$

Note that the term in the square brackets is the KL divergence between two distributions $\Pr(\tilde{y} | \mathbf{x}, \mathbf{w}_*)$ and $\Pr(\tilde{y} | \mathbf{x}, \mathbf{w})$ plus a constant independent of \mathbf{w} . Therefore $\mathbf{w} = \mathbf{w}_*$ minimizes this term and hence minimizes $L(\mathbf{w})$.

2.2 Proof of Lemma 1

We prove the strong convexity parameter here.

$$\begin{aligned} Q(\theta) &= \frac{1}{2} \int_{\eta(\mathbf{x}) \geq \theta} (\theta^2 - 2\eta(\mathbf{x})\theta + \eta(\mathbf{x})^2) d\mu(\mathbf{x}) + \frac{1}{2} \pi \theta^2 \\ &= \frac{1}{2} \theta^2 (\rho_\theta + \pi) - \theta \int_{\eta(\mathbf{x}) \geq \theta} \eta(\mathbf{x}) d\mu(\mathbf{x}) + c \end{aligned}$$

where $\rho_\theta = \int_{\eta(\mathbf{x}) \geq \theta} d\mu(\mathbf{x})$, c is a constant independent of θ . Then we can see the strong convexity parameter of $Q(\theta)$ over $[0, 0.5]$ is $\pi + \min_{\theta \in [0, 0.5]} \rho_\theta$.

2.3 Proof of Lemma 2

Proof. For $A \subseteq \mathcal{X}$, define $\rho(A) = \int_{\mathbf{x} \in A} 1 \cdot d\mu(\mathbf{x}) = \Pr(\mathbf{x} \in A)$. Let $\mathcal{X}_* = \{\mathbf{x} \in \mathcal{X} | \eta(\mathbf{x}) \geq \theta_*\}$ and $\mathcal{X}' = \{\mathbf{x} \in \mathcal{X} | \eta(\mathbf{x}) \geq \theta\}$, and note that $\eta_\theta(\mathbf{x}) = \mathbb{I}(\eta(\mathbf{x}) \geq \theta)$, we have

$$\frac{1}{2} F(\eta_\theta) = \frac{\int_{\mathcal{X}'} \eta(\mathbf{x}) d\mu(\mathbf{x})}{\pi + \rho(\mathcal{X}')} \quad (8)$$

According to [3], $F(\eta_{\theta_*}) = 2\theta_*$. Thus,

$$\begin{aligned} \theta_* &= \frac{1}{2} F(\eta_{\theta_*}) = \frac{\int_{\mathcal{X}_*} \eta(\mathbf{x}) d\mu(\mathbf{x})}{\pi + \rho(\mathcal{X}_*)} \\ \int_{\mathcal{X}_*} \eta(\mathbf{x}) d\mu(\mathbf{x}) &= \theta_* (\pi + \rho(\mathcal{X}_*)) \end{aligned} \quad (9)$$

Then we consider two cases based on the relation between θ and θ_* .

Case 1. $0 \leq \theta \leq \theta_*$

Since $\mathcal{X}_* \subseteq \mathcal{X}'$, let $A = \mathcal{X}' - \mathcal{X}_* = \{\mathbf{x} \in \mathcal{X} | \theta \leq \eta(\mathbf{x}) < \theta_*\}$. From (8),

$$\frac{1}{2}F(\eta_\theta) = \frac{\int_{\mathcal{X}_*} \eta(\mathbf{x})d\mu(\mathbf{x}) + \int_A \eta(\mathbf{x})d\mu(\mathbf{x})}{\pi + \rho(\mathcal{X}_*) + \rho(A)}$$

On A , we have $\eta(\mathbf{x}) \geq \theta$, thus $\int_A \eta(\mathbf{x})d\mu(\mathbf{x}) \geq \theta\rho(A)$. From (9), we have

$$\begin{aligned} \frac{1}{2}F(\eta_\theta) &\geq \frac{\theta_*(\pi + \rho(\mathcal{X}_*)) + \theta\rho(A)}{\pi + \rho(\mathcal{X}_*) + \rho(A)} = \frac{\theta_*(\pi + \rho(\mathcal{X}_*)) + \theta_*\rho(A) - \theta_*\rho(A) + \theta\rho(A)}{\pi + \rho(\mathcal{X}_*) + \rho(A)} \\ &= \theta_* - \frac{(\theta_* - \theta)\rho(A)}{\pi + \rho(\mathcal{X}_*) + \rho(A)} \geq \theta_* - (\theta_* - \theta) = \theta \end{aligned}$$

Thus $F(\eta_{\theta_*}) - F(\eta_\theta) \leq 2(\theta_* - \theta) \leq \frac{2}{\pi}|\theta_* - \theta|$.

Case 2. $\theta_* < \theta \leq 0.5$

Since $\mathcal{X}' \subseteq \mathcal{X}_*$, let $A = \mathcal{X}_* - \mathcal{X}' = \{\mathbf{x} \in \mathcal{X} | \theta_* \leq \eta(\mathbf{x}) < \theta\}$. From (8),

$$\frac{1}{2}F(\eta_\theta) = \frac{\int_{\mathcal{X}_*} \eta(\mathbf{x})d\mu(\mathbf{x}) - \int_A \eta(\mathbf{x})d\mu(\mathbf{x})}{\pi + \rho(\mathcal{X}_*) - \rho(A)}$$

On A , we have $\eta(\mathbf{x}) < \theta$, thus $\int_A \eta(\mathbf{x})d\mu(\mathbf{x}) \leq \theta\rho(A)$. From (9), we have

$$\begin{aligned} \frac{1}{2}F(\eta_\theta) &\geq \frac{\theta_*(\pi + \rho(\mathcal{X}_*)) - \theta\rho(A)}{\pi + \rho(\mathcal{X}_*) - \rho(A)} = \frac{\theta_*(\pi + \rho(\mathcal{X}_*)) - \theta_*\rho(A) + \theta_*\rho(A) - \theta\rho(A)}{\pi + \rho(\mathcal{X}_*) - \rho(A)} \\ &= \theta_* - \frac{(\theta - \theta_*)\rho(A)}{\pi + \rho(\mathcal{X}_*) - \rho(A)} \geq \theta_* - \frac{1}{\pi}(\theta - \theta_*). \end{aligned}$$

The last step holds because $\rho(A) < 1$ and $\rho(\mathcal{X}_*) - \rho(A) = \rho(\mathcal{X}') \geq 0$. Then we have $F(\eta_{\theta_*}) - F(\eta_\theta) \leq 2(\theta_* - \theta_* + \frac{1}{\pi}(\theta - \theta_*)) = \frac{2}{\pi}(\theta - \theta_*) = \frac{2}{\pi}|\theta_* - \theta|$.

We combine both cases and get the final result. \square

2.4 Proof of Theorem 2

Proof. Here we consider any stage k . Let τ denote the iteration index of SFO and $t = T_0 + \tau$ denote the global index. Define $g(\theta) = q(\theta) = \partial Q(\theta)$, $\mathbf{z} = (\mathbf{x}, y)$, $G(\theta, \mathbf{z}) = \pi\theta - (\eta(\mathbf{x}) - \theta)_+$, $\hat{G}_t(\theta, \mathbf{z}) = \hat{\pi}_t\theta - (\hat{\eta}_t(\mathbf{x}) - \theta)_+$. It is clear that $\mathbb{E}[G(\theta, \mathbf{z})] = g(\theta)$, and $\max(|g(\theta_\tau)|, |G(\theta_\tau, \mathbf{z}_t)|, |\hat{G}_t(\theta_\tau, \mathbf{z}_t)|) \leq 2$ for any τ . Following standard analysis of gradient descent, we have

$$\frac{1}{T} \sum_{\tau=1}^T (\theta_\tau - \theta_*) \hat{G}_t(\theta_\tau, \mathbf{z}_t) \leq \frac{|\theta_1 - \theta_*|^2}{2\gamma T} + \frac{\gamma \max(\hat{G}_t(\theta_\tau, \mathbf{z}_t))^2}{2}$$

Then by the convexity of $Q(\theta)$, we have

$$\begin{aligned} Q(\bar{\theta}_T) - Q(\theta_*) &\leq \frac{\|\theta_1 - \theta_*\|_2^2}{2\gamma T} + \frac{4\gamma}{2} + \frac{\sum_{\tau=1}^T (\theta_\tau - \theta_*)(g(\theta_\tau) - G(\theta_\tau, \mathbf{z}_t))}{T} \\ &\quad + \frac{\sum_{\tau=1}^T (\theta_\tau - \theta_*)(G(\theta_\tau, \mathbf{z}_t) - \hat{G}_t(\theta_\tau, \mathbf{z}_t))}{T} \\ &= \mathbf{I} + \mathbf{II} + \mathbf{III} + \mathbf{IV} \end{aligned}$$

Now we try to bound the four terms respectively. Note that $\mathbf{I} \leq \frac{R^2}{2\gamma T}$, $\mathbf{II} \leq 2\gamma$. To bound the third term, we utilize the similar analysis of SGD (e.g. [4]). Define

$$\begin{aligned} \tilde{\theta}_1 &= \theta_1 \in [0, 0.5] \cap \mathcal{B}(\theta_1, R), \\ \tilde{\theta}_{\tau+1} &= \Pi_{[0, 0.5] \cap \mathcal{B}(\theta_1, R)}(\tilde{\theta}_\tau - \gamma(g(\theta_\tau) - G(\theta_\tau, \mathbf{z}_t))). \end{aligned}$$

Then we have

$$\begin{aligned} \sum_{\tau=1}^T \gamma(\tilde{\theta}_\tau - \theta_*)(g(\theta_\tau) - G(\theta_\tau, \mathbf{z}_t)) &\leq \frac{\|\tilde{\theta}_1 - \theta_*\|_2^2}{2} + \frac{1}{2} \sum_{\tau=1}^T \gamma^2 \|g(\theta_\tau) - G(\theta_\tau, \mathbf{z}_t)\|_2^2 \\ &\leq \frac{R^2}{2} + 8\gamma^2 T. \end{aligned} \quad (10)$$

Note that both θ_τ and $\tilde{\theta}_\tau$ are measurable with respect to $\mathcal{F}_{t-1} = \{\mathbf{z}_1, \dots, \mathbf{z}_{t-1}\}$, and $\{S_\tau : \gamma(\theta_\tau - \tilde{\theta}_\tau)(g(\theta_\tau) - G(\theta_\tau, \mathbf{z}_t)), \tau = 1, \dots, T\}$ is a martingale difference sequence, and for any τ we have $|\gamma(\theta_\tau - \tilde{\theta}_\tau)(g(\theta_\tau) - G(\theta_\tau, \mathbf{z}_t))| \leq 4\gamma\|\theta_\tau - \tilde{\theta}_\tau\|_2 \leq 4\gamma \times 2R = 8\gamma R$. Then by Azuma-Hoeffding's inequality, we have with probability at least $1 - \frac{\delta}{3}$,

$$\sum_{\tau=1}^T \gamma(\theta_\tau - \tilde{\theta}_\tau)(g(\theta_\tau) - G(\theta_\tau, \mathbf{z}_t)) \leq 8\gamma R \sqrt{2T \ln\left(\frac{3}{\delta}\right)}. \quad (11)$$

Adding (10) and (11) together suffices to show that with probability at least $1 - \frac{\delta}{3}$, we have

$$\text{III} \leq \frac{R^2}{2\gamma T} + 8\gamma + \frac{8R\sqrt{2\ln\left(\frac{3}{\delta}\right)}}{\sqrt{T}}.$$

Next we bound **IV** according to the Lemma 3 introduced later. By union bound, we have with probability at least $1 - \frac{\delta}{3}$, we have

$$\begin{aligned} \text{IV} &\leq \frac{1}{T} \sum_{\tau=1}^T \left(\sup_{\tau} (\|\theta_\tau - \theta_1\|_2 + \|\theta_1 - \theta_*\|_2) \cdot \sup_{\theta \in [0, 0.5], \mathbf{z} \in \mathcal{Z}} \|\widehat{G}_t(\theta, \mathbf{z}) - G(\theta, \mathbf{z})\|_2 \right) \\ &\leq \frac{2R \cdot (1 + C\kappa) \times \sum_{t=1}^T \sqrt{\frac{\ln(12T/\delta)}{t}}}{T} \leq \frac{4R(1 + C\kappa)\sqrt{\ln\left(\frac{12T}{\delta}\right)}}{\sqrt{T}}, \end{aligned}$$

where the last inequality holds since $\sum_{t=1}^T \frac{1}{\sqrt{t}} \leq 2\sqrt{T}$. Combining these inequalities together, we have with probability at least $1 - \delta$, we have

$$Q(\bar{\theta}_T) - Q(\theta_*) \leq \frac{R^2}{\gamma T} + 10\gamma + \frac{R(20 + 4C\kappa)\sqrt{\ln(12T/\delta)}}{\sqrt{T}}.$$

Choosing $\gamma = \frac{R}{\sqrt{10T}}$, we have

$$Q(\bar{\theta}_T) - Q(\theta_*) \leq \frac{(2\sqrt{10} + (20 + 4C\kappa)\sqrt{\ln(12T/\delta)}) R}{\sqrt{T}}.$$

□

Lemma 3. *With probability at least $1 - \delta$,*

$$\sup_{\theta \in [0, 0.5], \mathbf{z} \in \mathcal{Z}} \|\widehat{G}_t(\theta, \mathbf{z}) - G(\theta, \mathbf{z})\|_2 \leq (1 + C\kappa) \sqrt{\frac{\ln(4/\delta)}{t}}.$$

Proof. For any θ and any \mathbf{z} , the following argument holds. By Hoeffding's inequality, we have with probability at least $1 - \frac{\delta}{2}$,

$$|\widehat{\pi}_t - \pi| \leq \sqrt{\frac{\ln(4/\delta)}{2t}}.$$

By the Assumption 1, we have with probability at least $1 - \frac{\delta}{2}$,

$$|\widehat{\eta}_t(\mathbf{x}) - \eta(\mathbf{x})| \leq C\kappa \sqrt{\frac{\ln(4/\delta)}{t}}.$$

Note that $0 \leq \theta \leq 0.5$, and hence we know that with probability at least $1 - \delta$,

$$\text{LHS} \leq |\widehat{\pi}_t - \pi| \cdot \theta + |\widehat{\eta}_t(\mathbf{x}_t) - \eta(\mathbf{x}_t)| \leq (1 + C\kappa) \sqrt{\frac{\ln(4/\delta)}{t}}.$$

□

2.5 Proof of Theorem 3

Given Theorem 2, the proof of Theorem 3 follows similar as the analysis [1] by noting that the objective function $Q(\theta)$ is strongly convex which is a special case of uniformly convex. For completeness, we give a proof here.

Proof. Define

$$\begin{aligned}\bar{\delta} &= \frac{2\delta}{\log_2 n}, \quad a(n, \bar{\delta}) = \frac{2\sqrt{10} + (20 + 4C\kappa)\sqrt{\ln(12n/\bar{\delta})}}{\sqrt{n}}, \\ \mu_0 &= \frac{2a(n_0, \bar{\delta})}{R_0}, \quad \mu_k = 2^k \mu_0, \quad R_k = R_0/2^k\end{aligned}$$

where $k = 1, \dots, m$. Then we have $\mu_k R_k^2 = 2^{-k} \mu_0 R_0^2$.

By definition of m in Algorithm 1 (FOFO), when $n \geq 100$,

$$0 < \frac{1}{2} \log_2 \frac{2n}{\log_2 n} - 2 \leq m \leq \frac{1}{2} \log_2 \frac{2n}{\log_2 n} - 1 \leq \frac{1}{2} \log_2 n, \quad (12)$$

so we have

$$2^m \geq \frac{1}{4} \sqrt{\frac{2n}{\log_2 n}}. \quad (13)$$

Define $c = \sqrt{\frac{2}{\sigma}}$, and note that $Q(\theta)$ is σ -strongly convex, and hence $\|\theta - \theta^*\|_2 \leq c(Q(\theta) - Q(\theta^*))^{\frac{1}{2}}$, where θ^* is the closest point to θ in $[0, 0.5]$.

Without loss of generality, we assume $c^2 \geq \frac{R_0}{2}$, i.e., $\frac{1}{c^2} \leq \frac{2}{R_0}$. Now we prove that $\frac{2}{R_0} \leq \mu_m$. When $n \geq 100$, we have

$$\begin{aligned}\mu_m &= 2^m \mu_0 \\ &\geq \frac{1}{4} \sqrt{\frac{2n}{\log_2 n}} \frac{4}{R_0} \left(\frac{\sqrt{10}}{\sqrt{n_0}} + \frac{(10 + 2C\kappa)\sqrt{\ln(12n_0/\bar{\delta})}}{\sqrt{n_0}} \right) \\ &\geq \frac{2}{R_0} \cdot \frac{1}{2} \sqrt{\frac{2n}{\log_2 n}} \left(\frac{\sqrt{10}}{\sqrt{n_0}} + \frac{8\sqrt{\ln(6\log_2 n)}}{\sqrt{n_0}} \right) \\ &\geq \frac{2}{R_0} \sqrt{\frac{2n}{\log_2 n}} \sqrt{\frac{(8\sqrt{10})\sqrt{\ln(6\log_2 n)}}{n_0}} \\ &\geq \frac{2}{R_0} \cdot \sqrt{\frac{2n}{\log_2 n}} \sqrt{\frac{(8\sqrt{10})\sqrt{\ln(3\log_2 n)}}{\frac{n}{m}}} \\ &\geq \frac{2}{R_0} \cdot \sqrt{\frac{2n}{\log_2 n}} \sqrt{\frac{(8\sqrt{10})\sqrt{\ln(3\log_2 n)}}{\frac{n}{\frac{1}{2} \log_2 \frac{2n}{\log_2 n} - 2}}} \\ &= \frac{2}{R_0} \sqrt{(8\sqrt{10})\sqrt{\ln(3\log_2 n)}} \left(1 - \frac{\log_2 \log_2 n + 3}{\log_2 n} \right) \\ &\geq \frac{2}{R_0}.\end{aligned}$$

where the first inequality holds because of (13), the second inequality stems from the fact that $10 + 2C\kappa > 8$, $0 < \delta < 1$, $n_0 \geq 1$, and the definition of $\bar{\delta}$, the third inequality holds by employing $a + b \geq 2\sqrt{ab}$, the fourth inequality holds because $0 < n_0 = \lfloor n/m \rfloor \leq n/m$, the fifth inequality holds because of the lower bound of m in (12), and the last inequality holds since when $n \geq 100$, the function $(8\sqrt{10})\sqrt{\ln(3\log_2 n)} \left(1 - \frac{\log_2 \log_2 n + 3}{\log_2 n} \right)$ is monotonically increasing with respect to n , and hence is greater than 1. So $\frac{2}{R_0} \leq \mu_m$. Recall that $\frac{1}{c^2} \leq \frac{2}{R_0}$, and thus, $\frac{1}{c^2} \leq \mu_m$.

Given $\hat{\theta}_k$, denote $\hat{\theta}_k^*$ by the closest optimal solution to $\hat{\theta}_k$. We consider two cases.

Case 1. If $\frac{1}{c^2} \geq \mu_0$, then $\mu_0 \leq \frac{1}{c^2} \leq \mu_m$. So there exists a k^* such that $\mu_{k^*} \leq \frac{1}{c^2} \leq \mu_{k^*+1} = 2\mu_{k^*}$, where $0 \leq k^* < m$. To utilize this fact, we have the following lemma.

Lemma 4. Let k^* satisfy $\mu_{k^*} \leq \frac{1}{c^2} \leq 2\mu_{k^*}$. Then for any $1 \leq k \leq k^*$, there exists a Borel set $\mathcal{A}_k \subset \Omega$ of probability at least $1 - k\bar{\delta}$, such that for $\omega \in \mathcal{A}_k$, the points $\{\hat{\theta}_k\}_{k=1}^m$ generated by the Algorithm 1 satisfy

$$\|\hat{\theta}_{k-1} - \hat{\theta}_{k-1}^*\|_2 \leq R_{k-1} = 2^{-k+1}R_0, \quad (14)$$

$$Q(\hat{\theta}_k) - Q_* \leq \mu_k R_k^2 = 2^{-k} \mu_0 R_0^2. \quad (15)$$

Moreover, for $k > k^*$ there is a Borel set $\mathcal{C}_k \subset \Omega$ of probability at least $1 - (k - k^*)\bar{\delta}$ such that on \mathcal{C}_k , we have

$$Q(\hat{\theta}_k) - Q(\hat{\theta}_{k^*}) \leq \mu_{k^*} R_{k^*}^2. \quad (16)$$

Proof. We prove (14) and (15) by induction. Note that (14) holds for $k = 1$. Assume it is true for some $k > 1$ on \mathcal{A}_{k-1} . According to the Theorem 2, there exists a Borel set \mathcal{B}_k with $\Pr(\mathcal{B}_k) \geq 1 - \bar{\delta}$ such that

$$Q(\hat{\theta}_k) - Q_* \leq R_{k-1} a(n_0, \bar{\delta}) = \frac{1}{2} \mu_k 2^{-k} R_0 R_{k-1} = \mu_k R_k^2,$$

which is (15). By the inductive hypothesis, $\|\hat{\theta}_{k-1} - \hat{\theta}_{k-1}^*\|_2 \leq R_{k-1}$ on the set \mathcal{A}_{k-1} . Define $\mathcal{A}_k = \mathcal{A}_{k-1} \cap \mathcal{B}_k$. Note that

$$\Pr(\mathcal{A}_k) \geq \Pr(\mathcal{A}_{k-1}) + \Pr(\mathcal{B}_k) - 1 \geq 1 - k\bar{\delta},$$

and on \mathcal{A}_k , by the strong-convexity of $Q(\theta)$ and the definition of k^* , we have

$$\|\hat{\theta}_k - \hat{\theta}_k^*\|_2^2 \leq c^2 (Q(\hat{\theta}_k) - Q_*) \leq \frac{Q(\hat{\theta}_k) - Q_*}{\mu_{k^*}} \leq \frac{\mu_k R_k^2}{\mu_{k^*}} \leq R_k^2,$$

which is (14) for $k + 1$.

Now we prove (16). For $k > k^*$, one can apply the similar strategy as in Theorem 2. Specifically, at the k -th stage with $k > k^*$, employing the similar proof of Theorem 2 by substituting all θ_* to $\hat{\theta}_{k-1}$, the first term of RHS becomes zero and hence we get a tighter bound of $Q(\hat{\theta}_k) - Q(\hat{\theta}_{k-1})$, we here relax the bound to be $R_{k-1} a(n_0, \bar{\delta})$.

So there exists a Borel set \mathcal{B}_k with $\Pr(\mathcal{B}_k) \geq 1 - \bar{\delta}$ such that

$$Q(\hat{\theta}_k) - Q(\hat{\theta}_{k-1}) \leq R_{k-1} a(n_0, \bar{\delta}) = 2^{k^*-k} R_{k^*-1} a(n_0, \bar{\delta}) = 2^{k^*-k} \mu_{k^*} R_{k^*}^2 = \mu_k R_k^2,$$

which implies that on $\mathcal{C}_k = \cap_{j=k^*+1}^k \mathcal{B}_j$, we have

$$Q(\hat{\theta}_k) - Q(\hat{\theta}_{k^*}) = \sum_{j=k^*+1}^k \left(Q(\hat{\theta}_j) - Q(\hat{\theta}_{j-1}) \right) \leq \sum_{j=k^*+1}^k 2^{k^*-j} \mu_{k^*} R_{k^*}^2 \leq \mu_{k^*} R_{k^*}^2.$$

By union bound, we have $\Pr(\mathcal{C}_k) = \Pr(\cap_{j=k^*+1}^k \mathcal{B}_j) \geq 1 - (k - k^*)\bar{\delta}$. Here completes the proof. \square

Now we proceed the proof as follows. Note that $\mu_0 \leq \frac{1}{c^2} \leq \mu_m$. At the end of k^* -th stage, on the Borel set \mathcal{A}_{k^*} of probability at least $1 - k^*\bar{\delta}$, we have

$$Q(\hat{\theta}_{k^*}) - Q_* \leq \mu_{k^*} R_{k^*}^2.$$

Then on the Borel set $\mathcal{D}_m = \mathcal{C}_m \cap \mathcal{A}_{k^*} = (\cap_{j=k^*+1}^m \mathcal{B}_j) \cap \mathcal{A}_{k^*}$ with $\Pr(\mathcal{D}_m) \geq 1 - m\bar{\delta}$, we have

$$\begin{aligned} Q(\hat{\theta}_m) - Q_* &= Q(\hat{\theta}_m) - Q(\hat{\theta}_{k^*}) + (Q(\hat{\theta}_{k^*}) - Q_*) \leq 2\mu_{k^*} R_{k^*}^2 \leq 4\left(\frac{\mu_{k^*}}{c^2}\right) \mu_{k^*} R_{k^*}^2 \\ &= (4c \cdot a(n_0, \bar{\delta}))^2. \end{aligned}$$

By the definition of m and $\bar{\delta}$, and the fact that $m \leq \frac{1}{2} \log_2 n$, we have $m\bar{\delta} \leq \delta$. So $\Pr(\mathcal{D}_m) \geq 1 - \delta$.

Table 1: Offline Testing F-measure (bold numbers represent the best performance)

Datasets	FOFO	OFO	LR	STAMP	OMCSL
webspam	.9348 ± .0003	.9348 ± .0004	.9347 ± .0005	.9312 ± .0014	.9282 ± .0046
a9a	.6789 ± .0015	.6755 ± .0020	.6518 ± .0026	.6735 ± .0034	.6704 ± .0096
ijcnn1	.6412 ± .0020	.5776 ± .0039	.4441 ± .0040	.5987 ± .0328	.6050 ± .0225
w8a	.7159 ± .0118	.6695 ± .0134	.6621 ± .0222	.6706 ± .0289	.6627 ± .0370
covtype (2 vs o)	.7627 ± .0005	.7625 ± .0005	.7557 ± .0004	.7568 ± .0055	.7557 ± .0081
covtype (1 vs o)	.7090 ± .0004	.7082 ± .0002	.6770 ± .0010	.7039 ± .0047	.7000 ± .0093
cov (3 vs o)	.7277 ± .0009	.7257 ± .0005	.6914 ± .0039	.7213 ± .0050	.7210 ± .0050
covtype (7 vs o)	.6723 ± .0022	.6521 ± .0025	.6140 ± .0037	.6417 ± .0197	.6513 ± .0150
covtype (6 vs o)	.4468 ± .0015	.4251 ± .0014	.1258 ± .0072	.3971 ± .0516	.4237 ± .0142
covtype (5 vs o)	.2648 ± .0036	.2488 ± .0027	.0000 ± .0000	.2218 ± .0246	.2362 ± .0304
covtype (4 vs o)	.5512 ± .0035	.5228 ± .0083	.4123 ± .0130	.3682 ± .0724	.5139 ± .0256
Sensorless (1 vs o)	.7549 ± .0047	.6732 ± .0022	.4774 ± .0156	.6243 ± .1394	.5401 ± .2360
Sensorless (2 vs o)	.4698 ± .0178	.2388 ± .0083	.1667 ± .0000	.3284 ± .1485	.4689 ± .0330
Sensorless (3 vs o)	.2138 ± .0047	.2254 ± .0048	.1345 ± .0709	.1819 ± .0812	.1804 ± .0413
Sensorless (4 vs o)	.5895 ± .0055	.3117 ± .0102	.1360 ± .0717	.3778 ± .2152	.4530 ± .0813
Sensorless (5 vs o)	.3089 ± .0049	.2343 ± .0047	.1009 ± .0868	.2264 ± .1186	.1782 ± .1228
Sensorless (6 vs o)	.3607 ± .0062	.2789 ± .0078	.0993 ± .0854	.2772 ± .0702	.2266 ± .1503
Sensorless (7 vs o)	.9994 ± .0002	.9996 ± .0001	.9986 ± .0010	.9988 ± .0009	.9982 ± .0017
Sensorless (8 vs o)	.4085 ± .0017	.3158 ± .0047	.0496 ± .0799	.3185 ± .1159	.3484 ± .0583
Sensorless (9 vs o)	.2783 ± .0037	.2069 ± .0039	.1346 ± .0710	.1749 ± .1352	.1902 ± .1251
Sensorless (10 vs o)	.6025 ± .0080	.4897 ± .0113	.1659 ± .0000	.4089 ± .2345	.5170 ± .0566
Sensorless (11 vs o)	.9997 ± .0000	.9997 ± .0002	.9998 ± .0002	.9997 ± .0001	.9998 ± .0002
protein (1 vs o)	.5008 ± .0026	.5037 ± .0059	.4643 ± .0114	.4914 ± .0163	.4930 ± .0116
protein (2 vs o)	.6849 ± .0035	.6835 ± .0040	.6390 ± .0053	.6787 ± .0069	.6735 ± .0144
protein (0 vs o)	.7479 ± .0017	.7483 ± .0014	.7183 ± .0023	.7430 ± .0071	.7423 ± .0052

Case 2. If $\frac{1}{c^2} < \mu_0$, then on $\mathcal{A}_1 = \mathcal{B}_1$,

$$Q(\hat{\theta}_1) - Q_* \leq R_0 \cdot a(n_0, \bar{\delta}) = \frac{R_0}{a(n_0, \bar{\delta})} \cdot a(n_0, \bar{\delta})^2 = \frac{2}{\mu_0} a(n_0, \bar{\delta})^2 \leq 2(c \cdot a(n_0, \bar{\delta}))^2.$$

Hence on $\mathcal{A}_1 \cap \mathcal{C}_m$, by using Lemma 4 and a similar argument as in case 1, we have

$$Q(\hat{\theta}_m) - Q_* = Q(\hat{\theta}_m) - Q(\hat{\theta}_1) + Q(\hat{\theta}_1) - Q_* \leq 2R_0 \cdot a(n_0, \bar{\delta}) \leq (2c \cdot a(n_0, \bar{\delta}))^2,$$

where $\Pr(\mathcal{A}_1 \cap \mathcal{C}_m) \geq 1 - \delta$.

Combining the two cases, we have with probability at least $1 - \delta$,

$$Q(\hat{\theta}_m) - Q_* \leq (4c \vee 2c)^2 (a(n_0, \bar{\delta}))^2 = \tilde{O}\left(\frac{\ln(\frac{1}{\delta})}{\sigma n}\right).$$

□

3 More Experimental Results

More experimental results are reported in Table 1 (offline testing results) and Figure 1 (online F-measure vs running time).

References

- [1] Anatoli Juditsky and Yuri Nesterov. Deterministic and stochastic primal-dual subgradient algorithms for uniformly convex minimization. *Stoch. Syst.*, 2014.
- [2] Nagarajan Natarajan, Oluwasanmi Koyejo, Pradeep Ravikumar, and Inderjit S. Dhillon. Consistent binary classification with generalized performance metrics. In *Neural Information Processing Systems (NIPS)*, 2014.




Figure 1: Online F-measure vs Running Time for more datasets

[3] Ming-Jie Zhao, Narayanan Edakunni, Adam Pocock, and Gavin Brown. Beyond fano's inequality: bounds on the optimal f-score, ber, and cost-sensitive risk and their implications. *Journal of Machine Learning Research*, 14(Apr):1033–1090, 2013.

[4] Martin Zinkevich. Online convex programming and generalized infinitesimal gradient ascent. 2003.