

Supplement to *A Polynomial-time Form of Robust Regression*

Yaliang Yu, Özlem Aslan, Dale Schuurmans

Department of Computing Science, University of Alberta, Edmonton AB T6G 2E8, Canada
{yaoliang, ozlem, dale}@cs.ualberta.ca

This supplement contains proofs of the claims made in the paper as well as additional technical developments. The supplement is organized into separate appendices to allow convenient spot checking of particular details of interest.

Appendix A Derivation details for the approximate estimation strategy.

Appendix B Robustness and non-robustness proofs.

Appendix C Consistency properties.

Appendix D Variational representation of loss functions
(including details on when such representations exist and how they can be recovered).

Appendix E Computational issues
(including notes on efficient implementation, and proofs of NP-hardness).

A Deriving the Approximate Estimator

Lemma 3.
$$\min_{0 \leq \eta \leq 1} \min_{\alpha} \boldsymbol{\eta}^T \ell(\mathbf{y} - K\alpha) + \mathbf{1}^T \boldsymbol{\psi}(\boldsymbol{\eta}) + \frac{\lambda}{2} \|\boldsymbol{\eta}\|_1 \alpha^T K \alpha \quad (24)$$

$$= \min_{0 \leq \eta \leq 1} \sup_{\boldsymbol{\nu}} \mathbf{1}^T \boldsymbol{\psi}(\boldsymbol{\eta}) - \boldsymbol{\eta}^T (\boldsymbol{\ell}^*(\boldsymbol{\nu}) - \Delta(\mathbf{y})\boldsymbol{\nu}) - \frac{1}{2\lambda} \boldsymbol{\nu}^T (K \circ (\boldsymbol{\eta} \|\boldsymbol{\eta}\|_1^{-1} \boldsymbol{\eta}^T)) \boldsymbol{\nu}. \quad (25)$$

where the function evaluations are componentwise.

Proof. The lemma amounts to dualizing the interior convex minimization problem

$$\min_{\alpha} \boldsymbol{\eta}^T \ell(\mathbf{y} - K\alpha) + \mathbf{1}^T \boldsymbol{\psi}(\boldsymbol{\eta}) + \frac{\lambda}{2} \|\boldsymbol{\eta}\|_1 \alpha^T K \alpha. \quad (26)$$

Recall from the main body that the function ℓ is assumed to be closed and convex. Therefore, strong Fenchel duality holds and we can reexpress ℓ as $\ell(r) = \sup_{\boldsymbol{\nu}} \boldsymbol{\nu}^T r - \boldsymbol{\nu}^T \boldsymbol{\ell}^*(\boldsymbol{\nu})$ where $\boldsymbol{\ell}^*$ is the Fenchel conjugate of ℓ [22, Ch.3]. Thus

$$\boldsymbol{\eta}^T \ell(\mathbf{y} - K\alpha) = \sup_{\boldsymbol{\nu}} \boldsymbol{\nu}^T \Delta(\boldsymbol{\eta})(\mathbf{y} - K\alpha) - \boldsymbol{\nu}^T \boldsymbol{\ell}^*(\boldsymbol{\nu}) \quad (27)$$

and we can therefore re-write (26) as

$$(26) = \min_{\alpha} \sup_{\boldsymbol{\nu}} \mathbf{1}^T \boldsymbol{\psi}(\boldsymbol{\eta}) - \boldsymbol{\eta}^T \boldsymbol{\ell}^*(\boldsymbol{\nu}) + \boldsymbol{\nu}^T \Delta(\mathbf{y})\boldsymbol{\eta} - \boldsymbol{\nu}^T \Delta(\boldsymbol{\eta})K\alpha + \frac{\lambda}{2} \|\boldsymbol{\eta}\|_1 \alpha^T K \alpha \quad (28)$$

$$= \sup_{\boldsymbol{\nu}} \min_{\alpha} \mathbf{1}^T \boldsymbol{\psi}(\boldsymbol{\eta}) - \boldsymbol{\eta}^T \boldsymbol{\ell}^*(\boldsymbol{\nu}) + \boldsymbol{\nu}^T \Delta(\mathbf{y})\boldsymbol{\eta} - \boldsymbol{\nu}^T \Delta(\boldsymbol{\eta})K\alpha + \frac{\lambda}{2} \|\boldsymbol{\eta}\|_1 \alpha^T K \alpha, \quad (29)$$

where (29) follows by strong duality (since for all fixed $\boldsymbol{\nu}$ in (29) the sublevel sets in α are bounded [22, Ch.3]).

Finally, α can be eliminated from the inner problem by solving for a critical point (the inner objective is convex in α). Taking the gradient in α and setting to zero gives the system of equations $\nabla_{\alpha} = K\Delta(\boldsymbol{\eta})\boldsymbol{\nu} + \lambda\|\boldsymbol{\eta}\|_1 K\alpha = 0$, which is satisfied by $\alpha = \Delta(\boldsymbol{\eta})\boldsymbol{\nu}/(\lambda\|\boldsymbol{\eta}\|_1)$. Substituting this solution back into (29) yields

$$(29) = \sup_{\boldsymbol{\nu}} \mathbf{1}^T \boldsymbol{\psi}(\boldsymbol{\eta}) - \boldsymbol{\eta}^T (\boldsymbol{\ell}^*(\boldsymbol{\nu}) - \Delta(\mathbf{y})\boldsymbol{\nu}) - \frac{1}{2\lambda\|\boldsymbol{\eta}\|_1} \boldsymbol{\nu}^T \Delta(\boldsymbol{\eta})K\Delta(\boldsymbol{\eta})\boldsymbol{\nu} \quad (30)$$

$$= \sup_{\boldsymbol{\nu}} \mathbf{1}^T \boldsymbol{\psi}(\boldsymbol{\eta}) - \boldsymbol{\eta}^T (\boldsymbol{\ell}^*(\boldsymbol{\nu}) - \Delta(\mathbf{y})\boldsymbol{\nu}) - \frac{1}{2\lambda} \boldsymbol{\nu}^T (K \circ (\boldsymbol{\eta} \|\boldsymbol{\eta}\|_1^{-1} \boldsymbol{\eta}^T)) \boldsymbol{\nu}, \quad (31)$$

establishing the lemma. ■

To prove the next lemma from the main body, recall the definitions.

$$\mathcal{N}_{\boldsymbol{\eta}} = \{N : N \succcurlyeq 0, N\mathbf{1} = \boldsymbol{\eta}, \text{rank}(N) = 1\} \quad (32)$$

$$\mathcal{M}_{\boldsymbol{\eta}} = \{M : M \succcurlyeq 0, M\mathbf{1} = \boldsymbol{\eta}, \text{tr}(M) \leq 1\}. \quad (33)$$

Lemma 4.

$$(25) = \min_{0 \leq \eta \leq 1} \min_{N \in \mathcal{N}_{\boldsymbol{\eta}}} \sup_{\boldsymbol{\nu}} \mathbf{1}^T \boldsymbol{\psi}(\boldsymbol{\eta}) - \boldsymbol{\eta}^T (\boldsymbol{\ell}^*(\boldsymbol{\nu}) - \Delta(\mathbf{y})\boldsymbol{\nu}) - \frac{1}{2\lambda} \boldsymbol{\nu}^T (K \circ N) \boldsymbol{\nu} \quad (34)$$

$$\geq \min_{0 \leq \eta \leq 1} \min_{M \in \mathcal{M}_{\boldsymbol{\eta}}} \sup_{\boldsymbol{\nu}} \mathbf{1}^T \boldsymbol{\psi}(\boldsymbol{\eta}) - \boldsymbol{\eta}^T (\boldsymbol{\ell}^*(\boldsymbol{\nu}) - \Delta(\mathbf{y})\boldsymbol{\nu}) - \frac{1}{2\lambda} \boldsymbol{\nu}^T (K \circ M) \boldsymbol{\nu}. \quad (35)$$

using the fact that $\mathcal{N}_{\boldsymbol{\eta}} \subseteq \mathcal{M}_{\boldsymbol{\eta}}$.

Proof. First, to prove the equality (34) consider any fixed $\boldsymbol{\eta} \in [0, 1]^n$. We will show that $N = \boldsymbol{\eta} \|\boldsymbol{\eta}\|_1^{-1} \boldsymbol{\eta}^T \Leftrightarrow N \in \mathcal{N}_{\boldsymbol{\eta}}$, which will immediately yield the equality. The direction \Rightarrow can be verified by a simple check. To prove \Leftarrow , assume $N \in \mathcal{N}_{\boldsymbol{\eta}}$. Since $N \succcurlyeq 0$ and $\text{rank}(N) = 1$ we know that $N = \mathbf{q}\lambda\mathbf{q}^T$ for some $\lambda > 0$ and \mathbf{q} such that $\|\mathbf{q}\| = 1$. But now, since $N\mathbf{1} = \boldsymbol{\eta}$, it must follow that $\mathbf{q} = \boldsymbol{\eta}/(\lambda\boldsymbol{\eta}^T\mathbf{1})$, hence $\mathbf{q} = \boldsymbol{\eta}/\|\boldsymbol{\eta}\|$ and $\lambda\mathbf{q}^T\mathbf{1} = \|\boldsymbol{\eta}\|$. Therefore, $\lambda = \|\boldsymbol{\eta}\|/\mathbf{q}^T\mathbf{1} = \|\boldsymbol{\eta}\|^2/(\boldsymbol{\eta}^T\mathbf{1})$. That is, N must have the form $N = \mathbf{q}\lambda\mathbf{q}^T = \boldsymbol{\eta}\|\boldsymbol{\eta}\|_1^{-1}\boldsymbol{\eta}^T$.

The inequality (35) then follows from the above argument, and the fact that $\|\boldsymbol{\eta}\|^2/(\boldsymbol{\eta}^T\mathbf{1}) \leq 1$, which implies $\text{tr}(N) \leq 1$ for any $N \in \mathcal{N}_{\boldsymbol{\eta}}$. Therefore $\mathcal{N}_{\boldsymbol{\eta}} \subseteq \mathcal{M}_{\boldsymbol{\eta}}$. ■

B Robustness

B.1 Non-robustness of (Non-constant) Convex Losses

Theorem 5. *Empirical risk minimization based on a (non-constant) convex loss cannot have bounded response if the domain (or kernel) is unbounded, even under Tikhonov regularization.*

We will separately prove the two cases; first, assuming an explicit feature representation expressed in a data matrix X , and second, in the case of an RKHS.

B.1.1 Explicit Feature Case

Proof. Consider the L_2 -norm regularized M -estimator:

$$\hat{\theta} = \arg \min_{\theta \in \Theta} \frac{\lambda}{2} \|\theta\|_2^2 + \sum_{i=1}^n \rho(y_i - X_i; \theta), \quad (36)$$

where ρ is some (non-constant) convex function. For simplicity, we assume ρ is differentiable (otherwise one can consider subdifferentials to arrive at the same conclusion).

Suppose the theorem is false, then $\hat{\theta}$ remains in a bounded interior subset. From the first order optimality condition (see, for instance, [22]), we know that

$$[\rho'(y_1 - X_1; \hat{\theta}) + \lambda]X_1 + \sum_{i=2}^n [\rho'(y_i - X_i; \hat{\theta}) + \lambda]X_i = 0. \quad (37)$$

Now we perturb (X_1, y_1) to cause a contradiction. Since ρ is a univariate convex and non-constant function, we apparently have $\lim_{d \rightarrow \infty} \rho'(d) > 0$ or $\lim_{d \rightarrow -\infty} \rho'(d) < 0$ or both. Assume $\lim_{d \rightarrow \infty} \rho'(d) > 0$ (the other case can be proved similarly).

Let y_1 and $\|X_1\|$ tend to infinity in a way that $\frac{y_1}{\|X_1\|}$ also tends to infinity. Then

$$y_1 - X_1; \hat{\theta} \geq y_1 - \|X_1\| \cdot \|\hat{\theta}\| \quad (38)$$

$$= \|X_1\| \cdot \left(\frac{y_1}{\|X_1\|} - \|\hat{\theta}\| \right) \quad (39)$$

tends to infinity since $\hat{\theta}$ is bounded. Therefore $\rho'(y_1 - X_1; \hat{\theta})$ tends to a positive number. But then the first term in (37) is unbounded in norm while the second term is bounded in norm (for we did not perturb $\{(X_i, y_i)\}_{i=2}^n$), contradiction. ■

Intuitively the meaning of this theorem is clear: If the loss function is unbounded, then any sample that is far enough can drag the estimate $\hat{\theta}$ outside of any bounded interior subset. Tikhonov regularization is not able to overcome this effect. Note that we need to perturb both X_1 and y_1 , in order to derive a contradiction (for instance, if one only perturbs y_1 , then convex functions with bounded derivatives will survive the proof).

B.1.2 RKHS Case

Proof. Let us consider the standard regularized M -estimator:

$$\hat{f}_{MR} \in \arg \min_{f \in \mathcal{H}} \frac{1}{n} \sum_{i=1}^n \rho(y_i - \langle \phi(x_i), f \rangle) + \frac{\lambda}{2} \|f\|_{\mathcal{H}}^2, \quad (40)$$

where $\{(x_i, y_i)\}_{i=1}^n$ are samples from the domain $\mathcal{X} \times \mathbb{R}$; and \mathcal{H} is the RKHS induced by some unbounded kernel κ , with its canonical feature map $\phi : \mathcal{X} \mapsto \mathcal{H}$. As above, we assume ρ is some non-constant convex loss function. For simplicity, we assume that the minimum in (40) is attained.

Suppose the theorem is false, then \hat{f}_{MR} remains in a bounded set. The first order necessary condition for the optimality of \hat{f}_{MR} yields

$$\rho'(y_1 - \langle \phi(x_1), \hat{f}_{MR} \rangle) \phi(x_1) + \sum_{i=2}^n \rho'(y_i - \langle \phi(x_i), \hat{f}_{MR} \rangle) \phi(x_i) + n\lambda \hat{f}_{MR} = 0, \quad (41)$$

where ρ' denotes the subdifferential of ρ (whose existence is guaranteed by convexity, moreover since ρ is finite-valued on \mathbb{R} , ρ' is also finite-valued on \mathbb{R}). The second term and third term above are bounded since \hat{f}_{MR} is assumed to be bounded. We will perturb (x_1, y_1) such that the first term is not bounded in norm, hence creating a contradiction.

Since ρ is convex and non-constant, we must have either $\lim_{y \rightarrow \infty} \rho'(y) > 0$ or $\lim_{y \rightarrow -\infty} \rho'(y) > 0$ (or both). Let us assume $\lim_{y \rightarrow \infty} \rho'(y) > 0$ (the other case can be proved similarly).

Let both y_1 and $\kappa(x_1, x_1)$ tend to infinity in a way such that $\frac{y_1}{\sqrt{\kappa(x_1, x_1)}}$ also tends to infinity. Then

$$y_1 - \langle \phi(x_1), \hat{f}_{MR} \rangle \geq y_1 - \sqrt{\kappa(x_1, x_1)} \|\hat{f}_{MR}\|_{\mathcal{H}} = \sqrt{\kappa(x_1, x_1)} \left(\frac{y_1}{\sqrt{\kappa(x_1, x_1)}} - \|\hat{f}_{MR}\|_{\mathcal{H}} \right), \quad (42)$$

hence

$$\left\| \rho' \left(y_1 - \langle \phi(x_1), \hat{f}_{MR} \rangle \right) \phi(x_1) \right\|_{\mathcal{H}} = \left| \rho' \left(y_1 - \langle \phi(x_1), \hat{f}_{MR} \rangle \right) \right| \sqrt{\kappa(x_1, x_1)} \rightarrow \infty, \quad (43)$$

due to our assumptions. ■

Note that again we need to perturb both $\kappa(x_1, x_1)$ and y_1 to reach a contradiction (for instance, if one only perturbs y_1 , then convex functions with bounded derivatives will survive our proof).

It should be clear in both these proofs that one may replace the L_2 -norm with other regularizers without affecting Theorem 5.

B.2 Robustness of the Approximate Regression Estimator

Theorem 3. Assume the loss ρ is bounded and has a variational representation (84) such that ℓ is Lipschitz-continuous and ψ' is bounded. Also assume there is at least one (unperturbed) inlier, and consider the perturbation of a single data point (y_1, x_1) . Under the following conditions, the rounded (re-optimized) estimator maintains bounded response:

- (i) If either y_1 remains bounded, or $\kappa(x_1, x_1)$ remains bounded.
- (ii) If $|y_1| \rightarrow \infty$, $\kappa(x_1, x_1) \rightarrow \infty$ and $\ell(y_1)/\kappa(x_1, x_1) \rightarrow \infty$.

To prove this theorem, we first need some key definitions. First recall the definition of outlier and inlier from the main body.

Definition 2 (Outliers and Inliers). For an L -Lipschitz loss ℓ , an outlier is a point (x_i, y_i) that satisfies $\ell(y_i) > L^2 K_{ii}/(2\lambda) - \psi'(0)$, while an inlier satisfies $\ell(y_i) + L^2 K_{ii}/(2\lambda) < -\psi'(1)$.

Also recall

$$\mathcal{N}_\eta = \{N : N \succcurlyeq 0, N\mathbf{1} = \eta, \text{rank}(N) = 1\} \quad (44)$$

$$\mathcal{M}_\eta = \{M : M \succcurlyeq 0, M\mathbf{1} = \eta, \text{tr}(M) \leq 1\}. \quad (45)$$

Let

$$R_0 = \min_{0 \leq \eta \leq 1} \min_{\alpha} \eta^T \ell(\mathbf{y} - K\alpha) + \mathbf{1}^T \psi(\eta) + \frac{\lambda}{2} \|\eta\|_1 \alpha^T K \alpha \quad (46)$$

$$R_1 = \min_{0 \leq \eta \leq 1} \min_{M \in \mathcal{M}_\eta} \sup_{\nu} \mathbf{1}^T \psi(\eta) + \eta^T (\Delta(\mathbf{y})\nu - \ell^*(\nu)) - \frac{1}{2\lambda} \nu^T (K \circ M) \nu \quad (47)$$

$$R_2 = \sup_{\nu} \mathbf{1}^T \psi(\eta) + \eta^T (\Delta(\mathbf{y})\nu - \ell^*(\nu)) - \frac{1}{2\lambda} \nu^T (K \circ (\eta \|\eta\|_1^{-1} \eta^T)) \nu \quad (48)$$

$$= \min_{\alpha} \eta^T \ell(\mathbf{y} - K\alpha) + \mathbf{1}^T \psi(\eta) + \frac{\lambda}{2} \|\eta\|_1 \alpha^T K \alpha, \quad (49)$$

where η in (48) is fixed at the optimal assignment determined by (47), and (49) follows by Lemma 3. Here, R_0 denotes the objective value obtained by the (intractable) oracle minimizer, R_1 is the objective value obtained by the relaxed solution, and R_2 denotes the objective value obtained by the rounded solution.

It is immediate that R_0 and R_1 must be bounded, since

$$R_1 \leq R_0 \leq n\psi(0) < \infty \quad (50)$$

where the first inequality follows from Lemma 4, and the second inequality is achieved by choosing $\eta = \alpha = 0$ in (47). The key question is whether R_2 , the rounded objective value, remains bounded. Once this is established for each of the cases, we finally show that this will imply that $\|\hat{f}_2\|_{\mathcal{H}}$ remains bounded, and the theorem will be proved.

B.2.1 Proof for Case (i), Bounded y

Proof. Assume that \mathbf{y} remains bounded. We will need to make use of the fact that, since ℓ is closed and convex, we have

$$\max_{\nu} \eta(y\nu - \ell^*(\nu)) = \eta \max_{\nu} y\nu - \ell^*(\nu) = \eta \ell(y). \quad (51)$$

Therefore, from (48) and (51) it follows that

$$R_2 \leq \sup_{\nu} \mathbf{1}^T \psi(\eta) - \eta^T (\ell^*(\nu) - \Delta(\mathbf{y})\nu) \quad (52)$$

$$= \mathbf{1}^T \psi(\eta) + \eta^T \ell(\mathbf{y}) \quad (53)$$

$$\leq n\gamma \quad (54)$$

such that $\gamma = \max_{0 \leq \eta \leq 1} \psi(\eta) + \max_{|y| \leq B} \ell(y) < \infty$. ■

B.2.2 Proof for Case (i), Bounded K

Proof. Assume K remains bounded; in particular that $|K_{ij}| \leq B$ for some $B < \infty$. We will need to make use of the fact that ℓ is Lipschitz-continuous. In particular, let ℓ have a Lipschitz-constant of L . Then it follows from Lemma 3 and the definition of Fenchel conjugate [26] that

$$R_2 = \sup_{-L \leq \nu \leq L} \mathbf{1}^T \psi(\boldsymbol{\eta}) - \boldsymbol{\eta}^T (\ell^*(\boldsymbol{\nu}) - \Delta(\mathbf{y})\boldsymbol{\nu}) - \frac{1}{2\lambda} \boldsymbol{\nu}^T (K \circ (\boldsymbol{\eta} \|\boldsymbol{\eta}\|_1^{-1} \boldsymbol{\eta}^T)) \boldsymbol{\nu} \quad (55)$$

$$\leq \sup_{-L \leq \nu \leq L} \mathbf{1}^T \psi(\boldsymbol{\eta}) - \boldsymbol{\eta}^T (\ell^*(\boldsymbol{\nu}) - \Delta(\mathbf{y})\boldsymbol{\nu}) \quad (56)$$

$$= \sup_{-L \leq \nu \leq L} \mathbf{1}^T \psi(\boldsymbol{\eta}) - \boldsymbol{\eta}^T (\ell^*(\boldsymbol{\nu}) - \Delta(\mathbf{y})\boldsymbol{\nu}) - \frac{1}{2\lambda} \boldsymbol{\nu}^T (K \circ M) \boldsymbol{\nu} + \frac{1}{2\lambda} \boldsymbol{\nu}^T (K \circ M) \boldsymbol{\nu} \quad (57)$$

$$\leq \sup_{-L \leq \nu \leq L} \left\{ \mathbf{1}^T \psi(\boldsymbol{\eta}) - \boldsymbol{\eta}^T (\ell^*(\boldsymbol{\nu}) - \Delta(\mathbf{y})\boldsymbol{\nu}) - \frac{1}{2\lambda} \boldsymbol{\nu}^T (K \circ M) \boldsymbol{\nu} \right\} + \sup_{-L \leq \nu \leq L} \frac{1}{2\lambda} \boldsymbol{\nu}^T (K \circ M) \boldsymbol{\nu} \quad (58)$$

$$= R_1 + \sup_{-L \leq \nu \leq L} \frac{1}{2\lambda} \boldsymbol{\nu}^T (K \circ M) \boldsymbol{\nu} \quad (59)$$

$$\leq n\psi(0) + \sup_{-L \leq \nu \leq L} \frac{1}{2\lambda} \boldsymbol{\nu}^T (K \circ M) \boldsymbol{\nu}. \quad (60)$$

Therefore it suffices to bound

$$\begin{aligned} & \sup_{-L \leq \nu \leq L} \boldsymbol{\nu}^T (K \circ M) \boldsymbol{\nu} \\ & \leq L^2 \text{tr}(KM) \end{aligned} \quad (61)$$

$$\leq L^2 Bn, \quad (62)$$

since $M \succcurlyeq 0$, $\text{tr}(M) \leq 1$, and $\lambda_{\max}(K) \leq Bn$ [25, p.292, Thm.5.6.9]. \blacksquare

B.2.3 Proof for Case (ii)

Proof. The proof in this case proceeds differently than the previous cases. Here we assume we are given a fixed data set $(x_1, y_1), \dots, (x_n, y_n)$, where only the first data point (x_1, y_1) is perturbed (without loss of generality), so that $|y_1| \rightarrow \infty$, $\kappa(x_1, x_1) \rightarrow \infty$ and $\ell(y_1)/\kappa(x_1, x_1) \rightarrow \infty$. Note that such a point must eventually become an outlier. We will show that this forces the corresponding weight η_1 to eventually satisfy $\eta_1 = 0$ in the relaxed solution (47), which will automatically imply that the rounded value R_2 stays at the same finite value thereafter (since no other data point is perturbed).

Consider the inner objective in (47):

$$\mathbf{1}^T \psi(\boldsymbol{\eta}) + \boldsymbol{\eta}^T (\Delta(\mathbf{y})\boldsymbol{\nu} - \ell^*(\boldsymbol{\nu})) - \frac{1}{2\lambda} \boldsymbol{\nu}^T (K \circ M) \boldsymbol{\nu} \quad (63)$$

$$= \mathbf{1}^T \psi(M\mathbf{1}) + \boldsymbol{\eta}^T (\Delta(\mathbf{y})\boldsymbol{\nu} - \ell^*(\boldsymbol{\nu})) - \frac{1}{2\lambda} \boldsymbol{\nu}^T (K \circ M) \boldsymbol{\nu}. \quad (64)$$

The gradients with respect to the parameters for this objective are given by

$$\nabla_{\boldsymbol{\nu}} = \Delta(\boldsymbol{\eta})(\mathbf{y} - \ell^*(\boldsymbol{\nu})') - \frac{1}{\lambda} (K \circ M) \boldsymbol{\nu} \quad (65)$$

$$\nabla_M = \psi'(M\mathbf{1})\mathbf{1}^T + (\Delta(\mathbf{y})\boldsymbol{\nu} - \ell^*(\boldsymbol{\nu}))\mathbf{1}^T - \frac{1}{2\lambda} (K \circ \boldsymbol{\nu}\boldsymbol{\nu}^T) \quad (66)$$

$$\frac{d}{d\eta_1} = \psi'(\eta_1) + y_1\nu_1 - \ell^*(\nu_1) - \frac{1}{2\lambda n} K_{1:} \boldsymbol{\nu}\nu_1. \quad (67)$$

Since ℓ is Lipschitz-continuous with Lipschitz constant L , we know that $|\nu| \leq L$ [26], hence

$$\left| \frac{1}{2\lambda n} K_{1:} \boldsymbol{\nu}\nu_1 \right| \leq \frac{\max |K_{1:}|}{2\lambda n} L^2 n = \frac{L^2}{2\lambda} K_{11}. \quad (68)$$

Now consider the tentative assignment $M_{1:} = \mathbf{0}^T$, $\eta_1 = 0$. At this assignment, all of the quadratic terms in ν_1 have been nullified in (63) and (64), leaving the optimization over ν_1 as

$$\max_{\nu_1} y_1 \nu_1 - \ell^*(\nu_1) = \ell(y_1) \quad (69)$$

by (51). Now note that at this solution for ν_1 we have

$$\frac{d}{d\eta_1} = \psi'(\eta_1) + \ell(y_1) - \frac{1}{2\lambda n} K_{1:} \boldsymbol{\nu} \nu_1 \quad (70)$$

$$\left. \frac{d}{d\eta_1} \right|_{\eta_1=0} = \psi'(0) + \ell(y_1) - \frac{1}{2\lambda n} K_{1:} \boldsymbol{\nu} \nu_1. \quad (71)$$

Therefore, if $\ell(y_1) > \frac{L^2}{2\lambda} K_{11} - \psi'(0)$ (the outlier condition) then $\left. \frac{d}{d\eta_1} \right|_{\eta_1=0} > 0$, which implies that η_1 stays at 0. We conclude that once the outlier condition is achieved (guaranteed by the assumptions), R_2 retains the same finite value after all subsequent perturbations of (x_1, y_1) , independent of $\ell(y_1)$ and $\kappa(x_1, x_1)$. ■

B.2.4 Final Step: Bounding $\|\hat{f}\|_{\mathcal{H}}$

Consider the rounded estimate \hat{f}_2 , corresponding to the solution to (49). It remains to bound $\|\hat{f}_2\|_{\mathcal{H}}$ in all the three cases discussed above.

Proof. Observe that

$$\|\hat{f}_2\|_{\mathcal{H}} = \boldsymbol{\alpha}^T K \boldsymbol{\alpha} \leq \frac{2R_2}{\lambda \|\boldsymbol{\eta}\|_1}. \quad (72)$$

Since R_2 has been proved bounded under the stated assumptions, we only need to consider the behavior of $\|\boldsymbol{\eta}\|_1$, which we do in two cases: $\boldsymbol{\eta} = 0$ and $\boldsymbol{\eta} \neq 0$.

First, assume $\boldsymbol{\eta} = 0$. Then $\boldsymbol{\alpha} = 0$ is an optimal solution of (49), implying that $\|\hat{f}_2\|_{\mathcal{H}} = 0$, which is obviously bounded.

So it only remains to consider $\boldsymbol{\eta} \neq 0$. In this case it suffices to bound $\|\boldsymbol{\eta}\|_1$ away from zero. We achieve this by appealing to the assumption that there is at least one *inlier*; that is, an unperturbed point (x_i, y_i) , $i \neq 1$, such that $\ell(y_i) + L^2 K_{ii}/(2\lambda) < -\psi'(1)$. For any such point, we can establish that $\eta_i = 1$, and the result will follow.

To show that any inlier gets weight η_i in the relaxed solution (47), tentatively consider the assignment $\eta_i = 1$ then recall from (67) that

$$\frac{d}{d\eta_i} = \psi'(\eta_i) + y_i \nu_i - \ell^*(\nu_i) - \frac{1}{2\lambda n} K_{i:} \boldsymbol{\nu} \nu_i \quad (73)$$

$$\left. \frac{d}{d\eta_i} \right|_{\eta_i=1} = \psi'(1) + y_i \nu_i - \ell^*(\nu_i) - \frac{1}{2\lambda n} K_{i:} \boldsymbol{\nu} \nu_i. \quad (74)$$

in the relaxed problem. By (69) above we know that $y_i \nu_i - \ell^*(\nu_i) \leq \ell(y_i)$. Furthermore, by (68) we know that $|K_{i:} \boldsymbol{\nu} \nu_i / (2\lambda n)| \leq L^2 K_{ii} / (2\lambda)$. Therefore, if the condition $\ell(y_i) + L^2 K_{ii} / (2\lambda) < -\psi'(1)$ is satisfied then $\left. \frac{d}{d\eta_i} \right|_{\eta_i=1} < 0$, which implies that η_i stays at 1. ■

C Consistency

C.1 Consistency of Standard Regression Estimators

Let us start by establishing some notation. Recall that \mathcal{H} is the RKHS induced by some kernel κ , $\{(x_i, y_i)\}_{i=1}^n$ are *i.i.d.* samples from the underlying training distribution $\mathbb{P}(x, y)$. For any function f (possibly random), define

$$\mathcal{R}(f) := \int_{\mathcal{X} \times \mathbb{R}} \rho(y - f(x)) d\mathbb{P}(x, y).$$

Then the following functions all aim at minimizing $\mathcal{R}(f)$ in one way or another:

$$f^* \in \underset{f}{\operatorname{argmin}} \mathcal{R}(f) \quad (75)$$

$$f_{\mathcal{H}} \in \underset{f \in \mathcal{H}}{\operatorname{argmin}} \mathcal{R}(f) \quad (76)$$

$$f_{\mathcal{H}, \lambda} \in \underset{f \in \mathcal{H}}{\operatorname{argmin}} \mathcal{R}(f) + \lambda \|f\|_{\mathcal{H}}^2 \quad (77)$$

$$\hat{f}_{\mathcal{H}, \lambda} \in \underset{f \in \mathcal{H}}{\operatorname{argmin}} \hat{\mathcal{R}}(f) + \lambda \|f\|_{\mathcal{H}}^2, \quad (78)$$

where $\hat{\mathcal{R}}(f) := \frac{1}{n} \sum_{i=1}^n \rho(y_i - f(x_i))$. For simplicity, we have tacitly assumed the existence of all minimizers in the above. We will also ignore measurability issues for the time being. Note that only the last function $\hat{f}_{\mathcal{H}, \lambda}$ depends on the data.

The regularized M -estimator $\hat{f}_{\mathcal{H}, \lambda}$ is said to be (loss) consistent if

$$\mathcal{R}(\hat{f}_{\mathcal{H}, \lambda}) - \mathcal{R}(f^*) \rightarrow 0 \quad (79)$$

as the sample size n increases to infinity and the regularization constant λ decreases to zero. To investigate when consistency can be assured, the following decomposition is standard and helpful [15]:

$$\mathcal{R}(\hat{f}_{\mathcal{H}, \lambda}) - \mathcal{R}(f^*) = \mathcal{R}(\hat{f}_{\mathcal{H}, \lambda}) - \mathcal{R}(f_{\mathcal{H}, \lambda}) - \lambda \|f_{\mathcal{H}, \lambda}\|_{\mathcal{H}}^2 \quad (80)$$

$$+ \mathcal{R}(f_{\mathcal{H}, \lambda}) + \lambda \|f_{\mathcal{H}, \lambda}\|_{\mathcal{H}}^2 - \mathcal{R}(f_{\mathcal{H}}) \quad (81)$$

$$+ \mathcal{R}(f_{\mathcal{H}}) - \mathcal{R}(f^*), \quad (82)$$

where the last term is usually called the approximation error, which measures how well can functions in \mathcal{H} approximate f^* under the ρ loss; the second term can be thought of as some stability error (where the regularization constant λ plays the role of perturbation); and the first term is related to the sampling error. Note that the last two terms depend only on the interaction between the RKHS \mathcal{H} (hence the kernel κ), the training distribution $\mathbb{P}(x, y)$, and the loss ρ . It is however independent of any estimation procedure (except perhaps providing some insights on how to practically tune the regularization constant). Very general bounds for the last two terms exist in the learning theory literature, see, for instance [15] and [27]. To make our presentation less complicated we will simply assume the sum of the last two terms, as a function of λ , goes to 0 when λ itself decreases to 0. The interested reader can consult the books [15] and [27] for precise technical conditions under which this is indeed so.

The right-hand side in (80) is apparently upper bounded by

$$\begin{aligned} \mathcal{R}(\hat{f}_{\mathcal{H}, \lambda}) + \lambda \|\hat{f}_{\mathcal{H}, \lambda}\|_{\mathcal{H}}^2 - \mathcal{R}(f_{\mathcal{H}, \lambda}) - \lambda \|f_{\mathcal{H}, \lambda}\|_{\mathcal{H}}^2 &= \mathcal{R}(\hat{f}_{\mathcal{H}, \lambda}) - \hat{\mathcal{R}}(\hat{f}_{\mathcal{H}, \lambda}) \\ &\quad + \hat{\mathcal{R}}(\hat{f}_{\mathcal{H}, \lambda}) + \lambda \|\hat{f}_{\mathcal{H}, \lambda}\|_{\mathcal{H}}^2 - \hat{\mathcal{R}}(f_{\mathcal{H}, \lambda}) - \lambda \|f_{\mathcal{H}, \lambda}\|_{\mathcal{H}}^2 \\ &\quad + \mathcal{R}(f_{\mathcal{H}, \lambda}) - \hat{\mathcal{R}}(f_{\mathcal{H}, \lambda}) \\ &\leq \mathcal{R}(\hat{f}_{\mathcal{H}, \lambda}) - \hat{\mathcal{R}}(\hat{f}_{\mathcal{H}, \lambda}) + \mathcal{R}(f_{\mathcal{H}, \lambda}) - \hat{\mathcal{R}}(f_{\mathcal{H}, \lambda}) \\ &\leq \sup_{f: \|f\|_{\mathcal{H}} \leq \frac{1}{\sqrt{\lambda}}} |\mathcal{R}(f) - \hat{\mathcal{R}}(f)|, \end{aligned} \quad (83)$$

where the first inequality is due to the optimality of $\hat{f}_{\mathcal{H}, \lambda}$, and the second inequality follows under the assumption $0 \leq \rho \leq 1$. Therefore we have related the right-hand side in (80) to the (uniform) sampling error. Applying standard uniform convergence bounds, for instance, the Rademacher complexity bound in [28, Theorem 8; Theorem 12; Lemma 22], leads to the following consistency result:

Proposition 1. *Assuming (81) and (82) approach 0 when $\lambda \rightarrow 0$, the loss ρ is Lipschitz and bounded between 0 and 1, $\sup_{x \in \mathcal{X}} \kappa(x, x) \leq 1$, then the regularized M -estimator defined in (78) is (ρ -loss) consistent.*

It is also possible to derive consistency results that depends on the capacity of the RKHS [15].

C.2 Consistency of Approximate Regression Estimators

Theorem 4. *Assume ℓ is Lipschitz-continuous and $\psi(\eta) = 1 - \eta$. Assume that the data is generated from a mixture of inliers and outliers, where $P(\text{inlier}) > P(\text{outlier})$. Then the estimate $\hat{\theta}$ produced by the rounded (re-optimized) method is loss consistent.*

Proof. From the proof of Theorem 3 above, we know that the relaxed solution will set $\eta_i = 0$ for all outliers, and $\eta_i = 1$ for all inliers. Since only outliers and inliers are present in the data, the solution η will be discrete $\{0, 1\}^n$, and all outliers will be ignored. Thus the reoptimized estimator \hat{f}_2 is based only on the inliers, and on such data the estimator is solved by a standard regularized empirical risk minimization. The consistency then follows from the standard results on regularized empirical risk minimization above. ■

D Variational Factorization of Loss Functions

Our objective in this section is to study some variational representation of loss functions. Our treatment is inspired by but different from that presented in [8].

Given any loss function $\rho(r)$ (not necessarily convex), we want to find (closed and proper) convex functions $\ell(r)$ and $\psi(\eta)$ such that

$$\rho(r) = \min_{\eta \in [0,1]} \eta \ell(r) + \psi(\eta). \quad (84)$$

Our main motivation to find such a variational representation for ρ is from optimization: the right hand side is separately (although not jointly) convex in both η and r , hence we can optimize ρ by dealing alternatively with ℓ and ψ , if the latter are in much simpler forms. This variational representation also allows us to develop a promising convex relaxation, as we will see soon.

Note that due to the closedness assumption on ψ , the minimum in (84) is always attained hence justifies our notation of \min instead of \inf . Also due to the convexity we pose on ℓ , the effective domain of ρ is necessarily convex. We also mention that the convexity of ψ simplifies our presentation but is not essential, for we can always replace ψ by its (closed) convex hull over the interval $[0, 1]$.

It is obvious that all closed convex functions ρ admit a variational representation (84): Just set $\ell(r) = \rho(r)$, $\psi(\eta) = \delta_{\{1\}}(\eta)$.¹² In fact, much more can be said. But let us prove a technical lemma first.

For any closed convex function $\psi(\eta)$, define its *restricted* Fenchel conjugate as:

$$\hat{\psi}(\hat{\eta}) := \max_{\eta \in [0,1]} \eta \hat{\eta} - \psi(\eta). \quad (85)$$

Note that $\hat{\psi}(\hat{\eta}) \leq \max_{\eta \in [0,1]} \eta \hat{\eta} - \min_{\eta \in [0,1]} \psi(\eta) = (\hat{\eta})_+ - \min_{\eta \in [0,1]} \psi(\eta) < \infty$, due to the closedness assumption on ψ . Hence $\hat{\psi}$ is defined on the whole real line.

Denote F as the set of all closed (proper) convex functions on $\bar{\mathbb{R}}$, \hat{F} the set of all closed (proper) convex functions $\hat{\ell}$ with effective domain \mathbb{R} and satisfying¹³

$$0 \leq \limsup_{\hat{\eta} \rightarrow -\infty} \frac{\hat{\psi}(\hat{\eta})}{\hat{\eta}} \leq \liminf_{\hat{\eta} \rightarrow \infty} \frac{\hat{\psi}(\hat{\eta})}{\hat{\eta}} \leq 1. \quad (86)$$

Define the mapping \mathcal{F} which maps $\psi \in F$ into its restricted Fenchel conjugate $\hat{\psi}$.

Lemma 5. \mathcal{F} maps F onto \hat{F} .

Proof. We first prove the range of the mapping \mathcal{F} is contained in \hat{F} . By definition,

$$\liminf_{\hat{\eta} \rightarrow \infty} \frac{\hat{\psi}(\hat{\eta})}{\hat{\eta}} = \liminf_{\hat{\eta} \rightarrow \infty} \max_{\eta \in [0,1]} \eta - \frac{\psi(\eta)}{\hat{\eta}} \quad (87)$$

$$\leq \liminf_{\hat{\eta} \rightarrow \infty} 1 - \frac{\min_{\eta \in [0,1]} \psi(\eta)}{\hat{\eta}} \quad (88)$$

$$= 1. \quad (89)$$

The last equality is because of the closedness of ψ . The other inequality in (86) can be similarly proved.

Let $\hat{\varphi} \in \hat{F}$, consider its (ordinary) Fenchel conjugate:

$$\hat{\varphi}^*(\eta) = \sup_{\hat{\eta}} \eta \hat{\eta} - \hat{\varphi}(\hat{\eta}) = \sup_{\hat{\eta}} \hat{\eta} \left(\eta - \frac{\hat{\varphi}(\hat{\eta})}{\hat{\eta}} \right). \quad (90)$$

¹² As is conventional in convex analysis, we use $\delta_C(\eta)$ to denote the indicator function for the point set C ; i.e., $\delta_C(\eta) = 0$ if $\eta \in C$, otherwise $\delta_C(\eta) = \infty$.

¹³ It does not matter if we change \limsup to \liminf or vice versa.

By the conditions (86) on \hat{F} , there exists a sequence $\hat{\eta}_i \rightarrow \infty$ as $i \rightarrow \infty$ such that $\frac{\hat{\psi}(\hat{\eta}_i)}{\hat{\eta}_i} \leq 1 + \epsilon$ for arbitrarily small $\epsilon > 0$ and all sufficiently large i . Now if $\eta > 1$ (hence $\eta \geq 1 + 2\epsilon$ for suitably small ϵ), then it is clear that $\hat{\varphi}^*(\eta) = \infty$. Similar arguments apply when $\eta < 0$, hence the effective domain of $\hat{\varphi}^*$ is contained in $[0, 1]$. Let $\psi = \hat{\varphi}^*$, then $\hat{\psi} = \hat{\varphi}^{**} = \hat{\varphi}$. This proves the mapping \mathcal{F} is onto. ■

Remark 1. *If we restrict the effective domain of the functions in F to be in $[0, 1]$, then an easy argument of duality establishes the one-one property of the mapping \mathcal{F} .*

Remark 2. *It should be clear that the interval $[0, 1]$ plays no special role here. Lemma 5 remains true (with obvious modifications) if we replace $[0, 1]$ with any other (non-degenerate) compact interval $[a, b]$. However, for our later convex relaxation, it is desirable to have $a \geq 0$.*

Theorem 5. *The function ρ admits a variational representation (84) if and only if $\rho = -\hat{\psi} \circ (-\ell)$ for some $\hat{\psi} \in \hat{F}$, $\ell \in F$. This shall be called the variational factorization of ρ .*

Proof. This is an immediate consequence of Lemma 5:

$$-\rho(r) := - \min_{\eta \in [0, 1]} \eta \ell(r) + \psi(\eta) \quad (91)$$

$$= \max_{\eta \in [0, 1]} \eta(-\ell(r)) - \psi(\eta) \quad (92)$$

$$= \hat{\psi}(-\ell(r)). \quad (93)$$

■

Let us now first illustrate a few examples to see how one can apply Theorem 5.

Example 1 (Variational representation of the Geman-McClure function $\frac{r^2}{1+r^2}$). *This example was also discussed in [8], through some integral arguments. Our derivation here, based on Theorem 5, is different. Rewrite $-\rho(r) = -1 + \frac{1}{1+r^2}$, which suggests us choose $\ell(r) = r^2$. Then $\hat{\psi}(\eta) = -1 + \frac{1}{1-\eta}$, $\forall \eta \leq 0$. Note that the decomposition $-\rho = \hat{\psi} \circ (-\ell)$ only determines $\hat{\psi}$ on the range of $-\ell$, hence giving us some flexibility to “complete” $\hat{\psi}$ on the entire real line such that (86) are satisfied. The easiest (nontrivial) way to extend $\hat{\psi}$ is to glue with it an affine function whose slope dominates $\hat{\psi}$. That is, define $\hat{\psi}(\eta) = a\eta + b$, $\forall \eta \geq 0$, where $a \geq \sup_{\eta \leq 0} \hat{\psi}'(\eta) = 1$. Picking $a = 1$ then satisfies both conditions in (86). To make $\hat{\psi}$ closed, we must set $b = \lim_{\eta \rightarrow 0} \hat{\psi}(\eta) = 0$. To summarize, $\rho(r) = \frac{r^2}{1+r^2}$ admits the variational representation (84) with $\ell(r) = r^2$, $\psi(\eta) = \hat{\psi}^*(\eta) = \sup_{\eta^*} \eta \eta^* - \hat{\psi}(\eta^*) = (\sqrt{\eta} - 1)^2$.*

Example 2 (Variational representation of the sigmoid function $\frac{\exp(r)}{1+\exp(r)}$). *This example is new. It was not discussed in [8], probably because there ℓ was restricted to the squared loss, which turns out to be prohibitive in this case. We simply repeat the procedure in Example 1. The similarity between the sigmoid function and the Geman-McClure function suggests us choose $\ell(r) = \exp(r)$, hence $\hat{\psi}(\eta)$ is accordingly determined on $(-\infty, 0)$ as $-1 + \frac{1}{1-\eta}$. Arguing exactly as in Example 1 we get the variational representation of the sigmoid function: $\ell(r) = \exp(r)$, $\psi(\eta) = (\sqrt{\eta} - 1)^2$.*

Example 3 (Variational representation of the Geman-Reynolds function $\frac{-1}{1+|r|}$). *This example was discussed in [8] where $\ell(r) = r^2$, but the final result turned out to be very complicated. We now demonstrate that by choosing a better ℓ , the representation can be significantly simplified. Note that $\rho(\eta) + 1 = \frac{|\eta|}{1+|\eta|}$. By analogy to the previous examples, it is easy to see that $\ell(r) = |r|$, $\psi(\eta) = (\sqrt{\eta} - 1)^2 - 1$ is a correct, and perhaps better, representation.*

The above three examples can be summarized as: $\frac{\ell(r)}{1+\ell(r)}$ with $\ell(r)$ positive and convex has variational representation $\ell(r)$, $\psi(\eta) = (\sqrt{\eta} - 1)^2$.

Example 4 (Some other examples). *These examples (with $\ell(r) = r^2$) were all discussed in [8]. We include them mainly for the purpose of completeness.*

Lorentzian function $\rho(r) = \log(1 + \ell(r))$ with ℓ positive and convex has variational representation: $\ell(r), \psi(\eta) = \eta - 1 - \log \eta$.

Tukey's biweight function $\rho(r) = \min\{\ell^2(r) - \ell^4(r) + \ell^6(r)/3, 1/3\}$ with ℓ convex has variational representation: $\ell(r), \psi(\eta) = \frac{1}{3} - \eta + \frac{2}{3}\eta^{3/2}$.

Leclerc's function $\rho(r) = 1 - \exp(-\ell^2(r))$ with ℓ convex has variational representation: $\ell(r), \psi(\eta) = \eta \log \eta - \eta + 1$.

The mean-field function $\rho(r) = -\log(1 + \exp(-\ell^2(r)))$ with variational representation: $\ell(r), \psi(\eta) = \eta \log \eta + (1 - \eta) \log(1 - \eta)$.

The nontrivial part in all examples is to identify the correct form of ℓ ; once that is done, the rest is fairly routine. Fortunately, for many useful loss functions, a few trial-and-error usually suffice. While we will develop a more principled way to yield a factorization, we nevertheless find the naive guessing method convenient and that is why we present it first.

We now turn to two immediate questions about Theorem 5:

- Is the variational factorization unique?
- Do all functions admit a variational factorization?

The first question is easily seen to be false since one can scale (or more generally, affinely transform) $\hat{\psi}$ and ℓ accordingly. The answer to the second question is also negative, for as we noted before, the effective domain of ρ needs to be convex. Moreover, by Alexandrov's theorem, $-\rho$ must have second derivative almost everywhere, in order to be a candidate of compositions of convex-concave functions. But it is well-known that there exist plenty of continuous and nowhere differentiable functions. Nevertheless, as noted before, all convex functions do admit a variational factorization (another trivial factorization would be $\hat{\psi}(\eta) = \eta, \ell = \rho$, a concrete example of the non-uniqueness). All concave functions whose negations are in \hat{F} also admit a variational factorization (again, a trivial factorization would be $\hat{\psi} = -\rho, \ell(r) = -r$).

To have a more satisfying (and involved) answer for the above two questions, we need to put some conditions on the loss function and its variational factors.¹⁴ We shall call a variational factorization $(\hat{\psi}, \ell)$ minimal if $\hat{\psi}$ is strictly increasing, and for any other factorization $(\hat{\varphi}, \xi)$, the function $\hat{\psi}^{-1} \circ \hat{\varphi}$ is convex. Now we can have a positive answer for the uniqueness question.

Lemma 6. *Minimal factorizations, if exist, are unique up to some (appropriate) affine transforms.*

Proof. Let $(\hat{\psi}, \ell), (\hat{\varphi}, \xi)$ both be minimal, then from the definition, we know both $\hat{\psi}^{-1} \circ \hat{\varphi}$ and $\hat{\varphi}^{-1} \circ \hat{\psi}$ are convex. Since both $\hat{\psi}$ and $\hat{\varphi}$ are assumed to be strictly increasing, it suffices to prove that only certain affine function can possibly be strictly increasing, convex, and having a convex inverse. But this is immediate, for we have $\forall \lambda \in [0, 1]$,

$$\lambda x + (1 - \lambda)y = f^{-1} \circ f(\lambda x + (1 - \lambda)y) \tag{94}$$

$$\leq f^{-1}(\lambda f(x) + (1 - \lambda)f(y)) \tag{95}$$

$$\leq \lambda f^{-1} \circ f(x) + (1 - \lambda)f^{-1} \circ f(y) \tag{96}$$

$$= \lambda x + (1 - \lambda)y. \tag{97}$$

The equality for all pairs of (x, y) forces f to be affine. ■

The next result requires some tools from real analysis, in particular, the Riesz representation theorem for bounded linear functionals (for a review, refer to [30]).

Lemma 7. *Let $-\rho$ be continuous and strictly decreasing on the open interval $\mathbb{I} \subseteq \mathbb{R}$ and admits a variational factorization $(\hat{\psi}, \ell)$ which are convex, continuous and strictly increasing on $\{-\ell(r) : r \in \mathbb{I}\}$ and \mathbb{I} , respectively, then*

1. *the right derivative, ρ_r , of ρ exists on \mathbb{I} ;*

¹⁴ Our treatment here is mainly inspired by [29].

2. $\log \rho_r$ is locally of bounded variation, hence corresponds (uniquely) to a (regular) Radon measure on \mathbb{I} , which we denote as $d(\log \rho_r)$;
3. a minimal factorization of $-\rho$ is given as $(\hat{\varphi}, \xi)$, with

$$\xi(x) := \int_a^x \exp\left(\int_b^y d\nu_+\right) dy, \quad \hat{\varphi}(x) = -\rho(\xi^{-1}(-x)), \quad (98)$$

provided that (98) is finite valued. Here $a = \inf \mathbb{I}$, $b \in \mathbb{I}$ and $d\nu_+$ is the positive part of the Radon measure $d(\log \rho_r)$.

Proof. 1. The proof is obvious hence omitted.

2. Due to the chain rule for derivatives:

$$(-\rho)_r = (\hat{\psi} \circ (-\ell))_r \quad (99)$$

$$= (\hat{\psi}_r \circ (-\ell)) \cdot (-\ell_r), \quad \text{hence} \quad (100)$$

$$\log \rho_r = \log(\hat{\psi}_r \circ (-\ell)) + \log \ell_r. \quad (101)$$

Note that all quantities inside the \log are positive due to the monotonicity assumption. Moreover, the two terms on the right hand side are decreasing and increasing, respectively, due to the convexity assumption. Therefore $\log \rho_r$ is locally of bounded variation. Through the usual Riemann-Stieltjes integral, $\log \rho_r$ induces a bounded linear functional on $C_c(\mathbb{I})$ (continuous functions on \mathbb{I} with compact support), hence by the Riesz representation theorem, corresponds to a unique (regular) Radon measure on \mathbb{I} .

3. Suppose $(\hat{\varphi}, \xi)$ is the minimal factorization and $(\hat{\psi}, \ell)$ is any variational factorization of $-\rho$, then from $-\rho = \hat{\varphi} \circ (-\xi) = \hat{\psi} \circ (-\ell)$ and the minimality of $\hat{\varphi}$ we know $f := \hat{\varphi}^{-1} \circ \hat{\psi} = (-\xi) \circ (-\ell)^{-1}$ is convex, hence $-\xi = f \circ (-\ell)$. It follows that $\log \xi_r = \log(f_r \circ (-\ell)) + \log \ell_r$. Since f is convex, $\log(f_r \circ (-\ell))$ is decreasing, therefore $d(\log \xi_r) \leq d(\log \ell_r)$. This proves some minimal property of the induced Radon measure of the minimal factorization.

Recall (101) which decomposes $d(\log \rho_r)$ into a positive measure and a negative one. The result above suggests us choose $d(\log \xi_r) = d\nu_+$, for the latter, being the positive part, is minimal by the Hahn-Jordan decomposition. This choice also forces $\hat{\varphi}_r$ to be increasing. Assuming $(\hat{\varphi}, \xi)$ as defined in (98) is finite valued, it is easy to verify the required convexity and monotonicity. \blacksquare

Remark 3. A careful inspection of the proof reveals that the minimal factorization in (98) does not hinge on the fact that $-\rho$ indeed admits a variational factorization, as long as $\log \rho_r$ is locally of bounded variation. Therefore finding the minimal factorization does not involve more work than finding merely a factorization.

Again, we digest the previous lemma through an example.

Example 5 (The sigmoid function revisited). $\mathbb{I} = (-\infty, \infty)$ in this example, hence $a = -\infty$ and we pick $b = 0$ (any other value would lead to the same result up to some constants). Simple calculation confirms $d(\log \rho_r) = \frac{1 - \exp(-x)}{1 + \exp(x)} dx$, where dx , as usual, is the Lebesgue measure on \mathbb{I} . Therefore $d\nu_+ = \mathbf{1}\{x \leq 0\} \cdot \frac{1 - \exp(-x)}{1 + \exp(x)} dx$. Direct integration in (98) gives $\xi(x) = \mathbf{1}\{x \leq 0\} \cdot \frac{4 \exp(x)}{1 + \exp(x)} + \mathbf{1}\{x > 0\} \cdot (2 + x)$ and $\hat{\varphi}(x) = \mathbf{1}\{x \geq -2\} \cdot \frac{x}{4} - \mathbf{1}\{x < -2\} \cdot \frac{1}{1 + \exp(x+2)}$. One easily verifies that the conditions in (86) are indeed satisfied. By Theorem 5 we know there exists a variational representation (φ, ξ) for the sigmoid function, although its explicit form is too cumbersome to write down. This variational factorization is very different from the one in Example 2, and perhaps is harder to come up with through guessing. From the point of view of optimization, the current factorization is inferior, despite of its theoretical value.

Lemma 7 cannot be applied directly to the Geman-McClure function due to the lack of monotonicity on the entire real line. However, such functions have the symmetry property that we can exploit. The idea is to restrict $\rho(x)$ to \mathbb{R}_+ where monotonicity is guaranteed, then apply Lemma 7 we get ξ on \mathbb{R}_+ (and $\hat{\varphi}$). Finally extend ξ to \mathbb{R}_- by symmetry.

Example 6 (The Geman-McClure function revisited). *Following the above recipe, $\mathbb{I} = (0, \infty)$, $a = 0$ and we leave b unspecified (as it only affects constants). Simple calculation confirms $d(\log \rho_r) = \frac{1-3x^2}{x(1+x^2)} dx$. Therefore $d\nu_+ = \mathbf{1}\{0 \leq x \leq 1/\sqrt{3}\} \cdot \frac{1-3x^2}{x(1+x^2)} dx$. Direct integration in (98) gives $\xi(x) = \mathbf{1}\{0 \leq x \leq 1/\sqrt{3}\} \cdot \frac{x^2}{1+x^2} + \mathbf{1}\{x > 1/\sqrt{3}\} \cdot (x + 1/4 - 1/\sqrt{3})$ and $\hat{\varphi}(x) = \mathbf{1}\{x \geq -1/4\} \cdot x - \mathbf{1}\{x < -1/4\} \cdot \frac{(x+1/4-1/\sqrt{3})^2}{1+(x+1/4-1/\sqrt{3})^2}$. For $x < 0$, we set $\xi(x) = \xi(-x)$ while leave $\hat{\varphi}$ untouched. One again verifies that the conditions in (86) are satisfied. By Theorem 5 we know there exists a variational representation (φ, ξ) for the sigmoid function, although its explicit form is again too cumbersome to write down. This variational factorization is very different from the one in Example 1, and is inferior from the point of view of optimization.*

The above two examples might lead one to conclude that the minimal factorization given by Lemma 7 is of little use from the viewpoint of optimization, which is our main motivation of developing the variational representation. However, as we point out below, the true value of Lemma 7 is to ensure one when a particular loss function does admit a variational factorization. Note that once the existence issue is solved, finding a meaningful factorization can usually be done through a few trial-and-error. We also notice that without the help of Lemma 7, it seems hard to envision the following theorem merely from the conclusion we drew in Theorem 5.

Theorem 6. *All bounded strictly increasing C^2 functions admit a variational factorization that satisfies (86), provided $\xi(x)$ defined below in (102) is finite valued and $\frac{\rho'(x)}{\xi'(x)}$ is bounded from above.*

Proof. Let ρ be bounded and strictly increasing. (98) now simplifies to

$$\xi(x) = \int_a^x \exp\left(\int_b^y \frac{\max\{\rho''(z), 0\}}{\rho'(z)} dz\right) dy. \quad (102)$$

We need only verify (86). Notice that $\hat{\varphi}$ in (98) is only determined in $\{-\xi(x) : x \in \text{dom}\rho\}$, but we can always complete it using the trick presented in Example 1 and 2.

Suppose $\hat{\varphi}(x)$ is defined for all sufficiently small x before we do the ‘‘completion’’, then

$$\limsup_{x \rightarrow -\infty} \frac{\hat{\varphi}(x)}{x} = \limsup_{x \rightarrow -\infty} \frac{-\rho(\xi^{-1}(-x))}{x} = 0, \quad (103)$$

since ρ is bounded. The other condition in (86) can be similarly argued. On the other hand, if $\hat{\varphi}(x)$ (for sufficiently small x) is only defined after we do the ‘‘completion’’, then we need to verify that $[-\rho(\xi^{-1}(-x))]' = \frac{\rho'(\xi^{-1}(-x))}{\xi'(\xi^{-1}(-x))} = \frac{\rho'(y)}{\xi'(y)}$ is between 0 and some positive number c (scaling $\hat{\varphi}$ and ξ appropriately will make $c = 1$). This holds because both ρ and ξ are strictly increasing, ρ' is bounded from above (for ρ is bounded), plus our assumption. ■

While it is possible to pin down a sufficient condition to remove the assumption in Theorem 6, we prefer not to do so, simply because this assumption itself is easily checkable. Apparently, Theorem 6 can be slightly modified to accommodate symmetric (and strictly increasing on, say, \mathbb{R}_+) loss functions. Refer to Examples 5 and 6.

Remark 4. *It is clear that we may replace the strictly decreasing assumption on the loss ρ with strictly increasing in both Lemma 7 and Theorem 6 (under obvious modifications), although such increasing functions seem to be uninteresting for loss minimization.*

In summary, we wonder what kind of functions can be factorized into the composition of a concave function and a convex function, where the concave part satisfies some condition like (86)? Of course, we would also like a principled method to find such factorizations, hopefully useful ones in the sense of optimization. Our results in Lemma 7 and Theorem 6 provide partial solutions to the above request, but seem to rely heavily on the monotonicity assumption. It would be nice to replace this assumption with something lighter.

E Computation

E.1 Efficient Implementation of the Approximate Estimator

Recall that the approximate estimator is computed in three steps: First, the relaxed lower bound (35) is computed, recovering $\boldsymbol{\eta}$ (see Lemma 4 in Appendix A). Second, the parameters $\boldsymbol{\alpha}$ are recovered by fixing $\boldsymbol{\eta}$ and re-solving for $\boldsymbol{\alpha}$ in (24) (see Lemma 3 in Appendix A). Third, $\boldsymbol{\eta}$ and $\boldsymbol{\alpha}$ are locally reoptimized in (24) using the previous $(\boldsymbol{\eta}, \boldsymbol{\alpha})$ as the initial point. We discuss an efficient implementation strategy, and the conditions under which polynomial run-time can be ensured.

For clarity we consider the case $\psi(\boldsymbol{\eta}) = 1 - \boldsymbol{\eta}$. (An efficient algorithm is possible for general ψ , but the details unnecessarily complicate the presentation.)

Relaxation: The first step of the approximate estimator is to compute the lower bound (35) given in Lemma 4, and recover the optimal $\boldsymbol{\eta}$ and M . This optimization can be solved efficiently as follows. Recall the definition $M_{\boldsymbol{\eta}} = \{M \succcurlyeq 0, M\mathbf{1} = \boldsymbol{\eta}, \text{tr}(M) \leq 1\}$, apply the definition $\psi(\boldsymbol{\eta}) = 1 - \boldsymbol{\eta}$, and observe

$$(35) = \max_{\boldsymbol{\nu}} \min_{0 \leq \boldsymbol{\eta} \leq \mathbf{1}} \min_{M \in M_{\boldsymbol{\eta}}} n - \mathbf{1}^T \boldsymbol{\eta} - \boldsymbol{\eta}^T (\boldsymbol{\ell}^*(\boldsymbol{\nu}) - \Delta(\mathbf{y})\boldsymbol{\nu}) - \frac{1}{2\lambda} \boldsymbol{\nu}^T (K \circ M) \boldsymbol{\nu} \quad (104)$$

$$= \max_{\boldsymbol{\nu}} \max_{\mathbf{a} \geq 0, \mathbf{b} \geq 0} \min_{\boldsymbol{\eta}} \min_{M \in M_{\boldsymbol{\eta}}} n - \mathbf{1}^T \boldsymbol{\eta} - \boldsymbol{\eta}^T (\boldsymbol{\ell}^*(\boldsymbol{\nu}) - \Delta(\mathbf{y})\boldsymbol{\nu}) - \frac{1}{2\lambda} \boldsymbol{\nu}^T (K \circ M) \boldsymbol{\nu} - \mathbf{a}^T M \mathbf{1} + \mathbf{b}^T (M \mathbf{1} - \mathbf{1}) \quad (105)$$

$$= \max_{\boldsymbol{\nu}} \max_{\mathbf{a} \geq 0, \mathbf{b} \geq 0} \min_{M \succcurlyeq 0, \text{tr}(M) \leq 1} n - \mathbf{1}^T M \mathbf{1} - \mathbf{1}^T M (\boldsymbol{\ell}^*(\boldsymbol{\nu}) - \Delta(\mathbf{y})\boldsymbol{\nu}) - \frac{1}{2\lambda} \boldsymbol{\nu}^T (K \circ M) \boldsymbol{\nu} - \mathbf{a}^T M \mathbf{1} + \mathbf{b}^T (M \mathbf{1} - \mathbf{1}) \quad (106)$$

$$= \max_{\boldsymbol{\nu}, \mathbf{a} \geq 0, \mathbf{b} \geq 0} n - \mathbf{b}^T \mathbf{1} - \max_{M \succcurlyeq 0, \text{tr}(M) \leq 1} \mathbf{1}^T M \mathbf{1} + \mathbf{1}^T M (\boldsymbol{\ell}^*(\boldsymbol{\nu}) - \Delta(\mathbf{y})\boldsymbol{\nu}) + \frac{1}{2\lambda} \boldsymbol{\nu}^T (K \circ M) \boldsymbol{\nu} + \mathbf{a}^T M \mathbf{1} - \mathbf{b}^T M \mathbf{1}, \quad (107)$$

where (104) follows by Sion's minimax theorem [26, Cor.37.3.2]; (105) follows by Lagrange duality; (106) follows by substituting $M\mathbf{1} = \boldsymbol{\eta}$ to replace the constraint and eliminate $\boldsymbol{\eta}$; and (107) is simple regrouping. Therefore (107) can be solved as a nonsmooth maximization:

$$\max_{\boldsymbol{\nu}, \mathbf{a} \geq 0, \mathbf{b} \geq 0} n - \mathbf{b}^T \mathbf{1} - f(\boldsymbol{\nu}, \mathbf{a}, \mathbf{b}), \quad (108)$$

where

$$f(\boldsymbol{\nu}, \mathbf{a}, \mathbf{b}) = \max_{M \succcurlyeq 0, \text{tr}(M) \leq 1} \mathbf{1}^T M \mathbf{1} + \mathbf{1}^T M \mathbf{c} + \frac{1}{2\lambda} \boldsymbol{\nu}^T (K \circ M) \boldsymbol{\nu} + \mathbf{a}^T M \mathbf{1} - \mathbf{b}^T M \mathbf{1} \quad (109)$$

$$= \frac{1}{2} \max_{M \succcurlyeq 0, \text{tr}(M) \leq 1} \text{tr}(MC(\boldsymbol{\nu}, \mathbf{a}, \mathbf{b})), \quad (110)$$

such that

$$C(\boldsymbol{\nu}, \mathbf{a}, \mathbf{b}) = 2\mathbf{1}\mathbf{1}^T + \mathbf{1}\mathbf{c}(\boldsymbol{\nu})^T + \mathbf{c}(\boldsymbol{\nu})\mathbf{1}^T + \frac{1}{\lambda} (K \circ \boldsymbol{\nu}\boldsymbol{\nu}^T) + \mathbf{a}\mathbf{1}^T + \mathbf{1}\mathbf{a}^T - \mathbf{b}\mathbf{1}^T - \mathbf{1}\mathbf{b}^T \quad (111)$$

$$\mathbf{c}(\boldsymbol{\nu}) = \boldsymbol{\ell}^*(\boldsymbol{\nu}) - \Delta(\mathbf{y})\boldsymbol{\nu}. \quad (112)$$

Note that in the maximization problem (108), $f(\boldsymbol{\nu}, \mathbf{a}, \mathbf{b})$ must be convex, since it is a pointwise maximum of linear functions [22, Ch.3]; hence (108) is a concave maximization problem. Each evaluation of $f(\boldsymbol{\nu}, \mathbf{a}, \mathbf{b})$ requires no more than $O(n^3)$ time, since (110) can be solved by computing the maximum eigenvector of $C(\boldsymbol{\nu}, \mathbf{a}, \mathbf{b})$ [31]. Moreover, a subgradient in $(\boldsymbol{\nu}, \mathbf{a}, \mathbf{b})$ can easily be recovered from the maximizer M , based on the fact that

$$\partial f \ni \begin{bmatrix} \Delta(M\mathbf{1})(\boldsymbol{\ell}'(\boldsymbol{\nu}) - \mathbf{y}) + \frac{1}{\lambda}(K \circ M)\boldsymbol{\nu} \\ M\mathbf{1} \\ -M\mathbf{1} \end{bmatrix} \quad (113)$$

by Danskin's theorem [32, Ch.6]. (Note that the maximizer M might not be unique, so f is non-smooth at points where this occurs.) Finally, at a solution $(\boldsymbol{\nu}, \mathbf{a}, \mathbf{b}; M)$ the weight vector $\boldsymbol{\eta}$ is recovered via $\boldsymbol{\eta} = M\mathbf{1}$.

Therefore, computing the relaxed solution requires solving a nonsmooth concave maximization over $3n$ variables, where each function evaluation (and subgradient) can be computed in $O(n^3)$ time. An ellipsoid algorithm can therefore be used to solve (108) in polynomial-time [21] [33].

Rounding: The rounding procedure involves a simple, smooth convex minimization of $\boldsymbol{\alpha}$ in (24), where $\boldsymbol{\eta}$ is fixed from the relaxation step. This problem can be solved in polynomial-time provided only that the base loss ℓ (assumed convex) is also self-concordant [20].

Reoptimization: Finally, in the reoptimization step, both $\boldsymbol{\eta}$ and $\boldsymbol{\alpha}$ are jointly (and locally) optimized in (24), starting from the solution above. A local optimum can, once again, be recovered in polynomial-time adding only the assumption that ψ (in addition to being convex) is also self-concordant.

Therefore, the estimation procedure requires only polynomial-time under the stated assumptions.

E.2 Proofs of NP-hardness

Our goal is to prove that minimizing a bounded loss function is NP-hard in general. After establishing the preliminary definitions required to be sufficiently precise about the results, we first prove that a special case—*clipped-loss* minimization—is strongly NP-hard in Section E.2.1. Then the NP-hardness of bounded loss minimization can be easily established by a reduction from clipped-loss minimization in Section E.2.2.

E.2.1 Hardness of Clipped Loss Minimization

Definition 3 (Loss function). *A function $\ell : \mathbb{R} \rightarrow \mathbb{R}$ is a loss function if (i) $\ell(r) \geq 0$ for all r ; (ii) $\ell(0) = 0$; and (iii) $\ell(r)$ is nondecreasing in r away from $r = 0$; that is, $r_1 \leq r_2 \leq 0$ implies $\ell(r_1) \geq \ell(r_2)$, and $0 \leq r_1 \leq r_2$ implies $\ell(r_1) \leq \ell(r_2)$.*

Definition 4 (τ -minimal two-sided loss). *A loss function ℓ is a τ -minimal two-sided loss if there exists a finite $B_\tau > 0$ such that (i) $r \leq B_\tau$ implies $\ell(r) \geq \tau$; and (ii) $r \geq B_\tau$ implies $\ell(r) \geq \tau$.*

Definition 5 (β -bounded loss). *A loss function ρ is a β -bounded loss if $\rho(y - \hat{y}) \leq \beta$ for all y and \hat{y} .*

Definition 6 (weakly τ -minimal loss). *A loss function ℓ is a weakly τ -minimal loss if it is a $(\tau - \epsilon)$ -minimal loss for all $\epsilon > 0$.*

Definition 7 (Clipped loss minimization).

Instance: $t \times n$ matrix X of training data, $t \times 1$ vector \mathbf{y} of training labels, 1-minimal two-sided loss function ℓ , nonnegative threshold number c .

Question: Is there an $n \times 1$ vector $\boldsymbol{\theta}$ such that $\sum_{i=1}^t \min(1, \ell(y_i - X_{i \cdot} \boldsymbol{\theta})) \leq c$?

Definition 8 (Bounded loss minimization).

Instance: $t \times n$ matrix X of training data, $t \times 1$ vector \mathbf{y} of training labels, 1-bounded and weakly 1-minimal two-sided loss function ℓ , nonnegative threshold number b .

Question: Is there an $n \times 1$ vector $\boldsymbol{\theta}$ such that $\sum_{i=1}^t \ell(y_i - X_{i \cdot} \boldsymbol{\theta}) \leq b$?

Definition 9 (Maximum 2-satisfiability).

Instance: Set U of variables, collection C of clauses over U such that each clause $c \in C$ has $|c| = 2$, positive integer $K \leq |C|$.

Question: Is there a truth assignment for U that simultaneously satisfies at least K of the clauses in C ?

Note that MAX2SAT is known to be NP-complete in general [34]. It is solvable in polynomial-time if $K = |C|$ [34], but NP-hard to approximate within a multiplicative constant better than $21/22 = 0.95454$ [35].

Theorem 7. *Clipped loss minimization is strongly NP-hard.*

Proof. We transform MAX2SAT to clipped-loss minimization. Let (U, C, K) constitute an instance of MAX2SAT.

Let $\mathbf{1}_i$ denote a boolean vector with a single 1 in position i . We will also use a scale factor s that will be specified below (we will be able to choose a value for s that is polynomial in $|U|$).

Widget construction: For each variable in $u_j \in U$ associate a feature $X_{:j}$ with corresponding weight θ_j . For each clause $c \in C$ construct three training examples in the form (\mathbf{x}^\top, y) as follows:

- For $u_i \vee u_j$ clauses, construct three examples $(s\mathbf{1}_i^\top, s)$, $(s\mathbf{1}_j^\top, s)$ and $(s(\mathbf{1}_i + \mathbf{1}_j)^\top, s)$.
- For $u_i \vee \neg u_j$ clauses, construct three examples $(s\mathbf{1}_i^\top, s)$, $(s\mathbf{1}_j^\top, 0)$ and $(s(\mathbf{1}_i - \mathbf{1}_j)^\top, 0)$.
- For $\neg u_i \vee u_j$ clauses, construct three examples $(s\mathbf{1}_i^\top, 0)$, $(s\mathbf{1}_j^\top, s)$ and $(s(\mathbf{1}_i - \mathbf{1}_j)^\top, 0)$.
- For $\neg u_i \vee \neg u_j$ clauses, construct three examples $(s\mathbf{1}_i^\top, 0)$, $(s\mathbf{1}_j^\top, 0)$ and $(s(\mathbf{1}_i + \mathbf{1}_j)^\top, s)$.

To illustrate how this construction will work, consider a training example $(s\mathbf{1}_i^\top, s)$. For a weight vector $\boldsymbol{\theta}$ one will obtain a prediction $\hat{y} = s\mathbf{1}_i^\top \boldsymbol{\theta} = s\theta_i$, which is compared to the target value $y = s$.




Figure 1: Depiction of the error surface in (θ_i, θ_j) for the three training examples constructed for each type of clause (when $i \neq j$): a $u_i \vee u_j$ clause, a $u_i \vee \neg u_j$ clause, a $\neg u_i \vee u_j$ clause, and a $\neg u_i \vee \neg u_j$ clause. (The trough widths are controlled by the scale factor s .) Note that the minimum total loss of 1 can only be achieved by setting (θ_i, θ_j) to a satisfying assignment in each of the four cases.

Note that if $\theta_i = 1$, then the loss must be zero on this example, since $\hat{y} = s = y$. However, as θ_i moves away from 1 the loss must increase until it “saturates”. Let

$$\rho(y - \hat{y}) = \min(1, \ell(y - \hat{y})) \quad (114)$$

denote the “clipped” loss. Since ℓ is a 1-minimal two-sided loss there must exist a finite $B_1 > 0$ such that for any y , $\hat{y} \leq y - B_1$ implies $\rho(y - \hat{y}) = 1$ and $\hat{y} \geq y + B_1$ implies $\rho(y - \hat{y}) = 1$. Conversely, if $\rho(y - \hat{y}) < 1$ then it must follow that $y - B_1 < \hat{y} < y + B_1$. The role of the scale factor s , therefore, is to control the width of the “trough” where the losses remain strictly less than 1: Note that for any scale factor $s > 0$, if $\rho(sy - s\hat{y}) < 1$ then $sy - B_1 < s\hat{y} < sy + B_1$, which holds if and only if $y - B_1/s < \hat{y} < y + B_1/s$. Thus, the larger the choice of s , the narrower the trough where losses less than 1 can be achieved for a given training example.

Figure E.2.1 depicts the error surfaces created by the set of three training examples for each of the four clause types (for sufficiently large s) when the variables are distinct, $i \neq j$. Note that the minimum loss any weight vector can achieve on a clause (i.e., on its associated set of three training examples) is always 1, and this can only be achieved by assigning boolean weights that “satisfy” the clause.




Figure 2: Illustrating bins of width $\frac{B_1\sqrt{2}}{s}$ between $\frac{B_1(1+\sqrt{2})}{s}$ and $\frac{1}{2}$, and their mirror images between $\frac{1}{2}$ and $1 - \frac{B_1(1+\sqrt{2})}{s}$.




Figure 3: Illustrating how an empty (bin, mirror bin) pair can be used to define which component weight values are “nearly discrete” versus “close to $1/2$ ”.

Choosing the scale factor s : The scale factor s needs to be fixed to a sufficiently large value so that, for any weight vector θ , there exists a “gap” between 0 and $\frac{1}{2}$ (and a corresponding mirror gap between $\frac{1}{2}$ and 1) that contains no individual weight component θ_i . In particular, consider a partition of the interval $[\frac{B_1(1+\sqrt{2})}{s}, \frac{1}{2}]$ into disjoint bins of size $\frac{B_1\sqrt{2}}{s}$. There are at least $\lfloor (\frac{1}{2} - \frac{B_1(1+\sqrt{2})}{s}) / \frac{B_1\sqrt{2}}{s} \rfloor = \lfloor \frac{s}{B_12\sqrt{2}} - \frac{1}{\sqrt{2}} - 1 \rfloor$ such bins that fit entirely within the interval.¹⁵ So, by setting

$$s = B_12\sqrt{2}(|U| + 3) \quad (115)$$

it follows that there must be at least $|U| + 1$ disjoint bins of width $\frac{B_1\sqrt{2}}{s}$ within the interval $[\frac{B_1(1+\sqrt{2})}{s}, \frac{1}{2}]$, plus a mirror set of $|U| + 1$ disjoint bins within the interval $[\frac{1}{2}, 1 - \frac{B_1(1+\sqrt{2})}{s}]$; see Figure 2. Given that the weight vector only has $|U|$ components, by construction, we know that for any particular θ there must be at least one (bin, mirror bin) pair such that neither bin contains any θ_i value; see Figure 3. We will use these empty bins to define which θ_i values are considered to be “nearly discrete” versus “close to $1/2$ ”, as shown in Figure 3.

Main argument: We now proceed to show that there is always a boolean weight vector that is a global minimizer of the total clipped loss. Let

$$\rho(\theta) = \sum_{i=1}^t \rho(y_i - X_i \cdot \theta) \quad (116)$$

denote the total clipped loss obtained by a weight vector θ on the constructed set of training examples. Let $\theta^{(0)}$ be a global minimizer of (116). Then we know that there must be some (bin, mirror bin) pair that contains no component of $\theta^{(0)}$, which we can use to define “nearly discrete” versus “close to $1/2$ ” weight values (Figure 3). Then by Lemma 8 below we know there must be some

¹⁵ The reason for considering the interval $[\frac{B_1(1+\sqrt{2})}{s}, \frac{1}{2}]$ instead of $[0, \frac{1}{2}]$ is that we also need to ensure the empty bin does not intersect the parallelogram at the intersection between an axis-parallel and diagonal trough. For a given s this will be true for any bin that is at least $\frac{B_1(1+\sqrt{2})}{s}$ away from 0 or 1.




Figure 4: Illustrating the rounding scheme of Lemma 8 for a $u_i \vee u_j$ clause. (The scheme for the other three clause types depicted in Figure E.2.1 is isomorphic. Weight pairs (θ_i, θ_j) in the four *red* corner quadrants are rounded to the boolean corners. Weight pairs (θ_i, θ_j) in the four *blue* semi-rectangles on the edges are rounded to the line where the “nearly discrete” value becomes boolean. Weight pairs (θ_i, θ_j) in the *green* central square are not rounded. No weight pair occurs in the cross-hatch formed by the empty bins.

weight vector $\theta^{(1)}$ which contains only boolean or “close to $1/2$ ” values such that $\rho(\theta^{(1)}) \leq \rho(\theta^{(0)})$. Then by Lemma 9 below we know there must exist a pure boolean weight vector $\theta^{(2)}$ such that $\rho(\theta^{(2)}) \leq \rho(\theta^{(1)})$. Finally given a boolean assignment $\theta^{(2)}$, a corresponding truth assignment $\mathbf{u}^{(2)}$ can be directly recovered via the transformation

$$\begin{aligned} u_i = \text{true} &\Leftrightarrow \theta_i = 1 \\ u_i = \text{false} &\Leftrightarrow \theta_i = 0. \end{aligned} \quad (117)$$

Furthermore, for a boolean weight vector $\theta^{(2)}$ we have that

$$\begin{aligned} \rho(\theta^{(2)}) &= 3 \times |\{\text{clauses falsified by } \mathbf{u}^{(2)}\}| \\ &\quad + 1 \times |\{\text{clauses satisfied by } \mathbf{u}^{(2)}\}| \end{aligned} \quad (118)$$

$$= 3|C| - 2 \times |\{\text{clauses satisfied by } \mathbf{u}^{(2)}\}|. \quad (119)$$

It then follows that $|\{\text{clauses satisfied by } \mathbf{u}^{(2)}\}| \geq K$ if and only if $\rho(\theta^{(2)}) \leq 3|C| - 2K$. That is, the minimum clipped loss achieved can be used to decide whether K clauses can be satisfied in the original instance. ■

Lemma 8. *For any weight vector θ there is always a weight vector $\tilde{\theta}$ with only boolean components or components that are “close to $1/2$ ” such that $\rho(\tilde{\theta}) \leq \rho(\theta)$.*

Proof. Fix θ and use one of its empty bins to define which component values are “nearly discrete” versus “close to $1/2$ ”. Let $\tilde{\theta}$ be the vector obtained by rounding all “nearly discrete” values of θ to their nearest boolean value. We show that the total clipped loss obtained by $\tilde{\theta}$ cannot be greater than that of θ .

Consider Figure 4, which depicts the effect of rounding for a clause of the $u_i \vee u_j$ type (the other three cases are isomorphic). For any clause, there are two situations to consider: when both weights are rounded and when only one weight is rounded.

First, consider the case where both weights (θ_i, θ_j) are “nearly discrete” and hence both are rounded to their nearest boolean value. This corresponds to being in one of the *red* corner quadrants shown in Figure 4. By inspection, one can see that the boolean assignment within each quadrant achieves the minimum loss attainable within the quadrant. In particular, in the three “satisfying” quadrants




Figure 5: Possible weight pairs that can occur after rounding.

the boolean assignment is the strict minimizer of the loss, whereas in the “falsifying” quadrant all assignments achieve the same loss of 3. Therefore, the total loss cannot have increased on such a clause due to rounding.

Second, consider the case where one of the weights is “nearly discrete” and hence rounded, while the other is “close to $1/2$ ” and hence not rounded. This corresponds to being in one of the *blue* semi-rectangles on the edges shown in Figure 4. Once again, by inspection one can see that rounding the “nearly discrete” weight to its nearest boolean value cannot increase the loss obtained. In particular, rounding the “nearly discrete” variable yields the minimum loss assignment in the top and right semi-rectangles, while every assignment in the bottom and left semi-rectangles obtains the same loss of 3. This latter property is precisely what is achieved by the bin definitions: the empty bin width of $\frac{B_1\sqrt{2}}{s}$ is sufficient to ensure that the *blue* semi-rectangles do not intersect the diagonal trough. Therefore, the total loss cannot have increased on such a clause due to rounding.

In clauses where both weights are “close to $1/2$ ” (the *green* square in the middle) neither weight is rounded, hence the loss does not change. The result follows. ■

Lemma 9. *For any weight vector θ with only boolean components or components that are “close to $1/2$ ” there is always a boolean weight vector $\hat{\theta}$ such that $\rho(\hat{\theta}) \leq \rho(\theta)$.*

Proof. Consider Figure 5, which depicts the possible weight pairs (θ_i, θ_j) participating in a clause of type $u_i \vee u_j$. (The situation for any of the clause types is isomorphic.) The isolated *red* points are the only purely boolean values, the *blue* line segments indicate pairs where one value is boolean, and the *green* square shows the region where both weights are “close to $1/2$ ”. Observe that any weight pair with one or more “close to $1/2$ ” values (i.e., the union of the *blue* lines and the *green* square) has loss at least 2.

Given the weight vector θ , partition the clauses into three subsets: the set of clauses where both weight values are boolean (“closed clauses”), the set of clauses where one weight is boolean and the other is not (“mixed clauses”), and the set of clauses where both weights are “close to $1/2$ ” (“open clauses”). Let the set of “non-closed clauses” consist of the union of the “mixed clauses” and the “open clauses”. From above, we know that θ achieves a loss of at least 2 on each “non-closed clause”. Let c denote the loss per clause achieved by θ on the “closed clauses”. There are two cases to consider.

Case 1: ($c \geq 2$) In this case the overall loss per clause achieved by θ must be at least 2, which implies the total loss is $\rho(\theta) \geq 2|C|$. Therefore, by Lemma 10 below we know that there must exist

some assignment $\hat{\mathbf{u}}$ to the variables in U that satisfies at least $\frac{|C|}{2}$ of the clauses in C . Let $\hat{\boldsymbol{\theta}}$ denote the corresponding boolean weight vector, recovered via the translation (117). By (119) we know that

$$\rho(\hat{\boldsymbol{\theta}}) = 3|C| - 2 \times |\{\text{clauses satisfied by } \mathbf{u}^{(2)}\}| \quad (120)$$

$$\leq 2|C| \quad (121)$$

$$\leq \rho(\boldsymbol{\theta}). \quad (122)$$

Case 2: ($c < 2$) Let B denote the indices where $\boldsymbol{\theta}$ is boolean, and let N denote the indices where $\boldsymbol{\theta}$ is “close to $1/2$ ”. (Since $c < 2$ there must be at least one component θ_i that is boolean.) Fix the boolean components of $\boldsymbol{\theta}_B$, which in turn fixes the outcomes on the “closed clauses”. For any “mixed clause” let θ_i denote the weight that is boolean. We would like to preserve the assignment of θ_i . Therefore, temporarily replace such a mixed clause with a new singleton clause defined by substituting u_i with its partner u_j . Consider the new set of altered “mixed clauses”, in union with the set of “open clauses”. Denote this new set \tilde{C} . Note that \tilde{C} is defined only on the subset of variables U_N corresponding to weights in $\boldsymbol{\theta}$ that are “close to $1/2$ ”. By Lemma 10 below, there must be some assignment $\tilde{\mathbf{u}}_N$ that satisfies at least half of the clauses in \tilde{C} , hence there exists a corresponding boolean weight vector $\tilde{\boldsymbol{\theta}}_N$ that achieves a loss of at most $2|\tilde{C}|$ on \tilde{C} .

Now consider the boolean weight vector $\hat{\boldsymbol{\theta}}$ formed by conjoining $\boldsymbol{\theta}_B$ with $\tilde{\boldsymbol{\theta}}_N$. Note that for any “closed clause” $\boldsymbol{\theta}$ and $\hat{\boldsymbol{\theta}}$ behave identically and hence achieve the same loss. However, recall $\boldsymbol{\theta}$ achieves a loss of at least 2 on each “non-closed clause”, while by the above construction, $\hat{\boldsymbol{\theta}}$ achieves a loss of at most 2 per clause over the “non-closed clauses”. (In particular, even though $\hat{\mathbf{u}}_N$ was constructed by satisfying temporarily altered “mixed clauses” above, if it satisfies such a clause, then it must also satisfy the original “mixed clause”.) Hence $\rho(\hat{\boldsymbol{\theta}}) \leq \rho(\boldsymbol{\theta})$.

Since in each of the two cases we were able to identify a boolean weight vector $\hat{\boldsymbol{\theta}}$ that achieves a loss no worse than $\boldsymbol{\theta}$, the result must follow. ■

Lemma 10. *For any MAX2SAT instance (U, C, K) and any assignment \mathbf{u} to its variables U , either \mathbf{u} or its negation $\neg\mathbf{u}$ must satisfy at least $\frac{|C|}{2}$ of the clauses in C .*

Proof. Consider any assignment \mathbf{u} . Note that $\neg\mathbf{u}$ must satisfy each clause that \mathbf{u} falsifies. Therefore, if \mathbf{u} satisfies fewer than $\frac{|C|}{2}$ clauses, it must falsify at least $\frac{|C|}{2}$ of the clauses, hence $\neg\mathbf{u}$ would have to satisfy at least $\frac{|C|}{2}$ of the clauses. Otherwise, \mathbf{u} satisfies at least $\frac{|C|}{2}$ of the clauses. ■

E.2.2 Hardness of Bounded Loss Minimization

Finally, we are in a position to prove Theorem 2 from the main body. Recall the definition of bounded loss minimization (Definition 8).

Theorem 2. *Bounded (non-constant) loss minimization is NP-hard.*

Proof. We transform MAX2SAT to bounded-loss minimization. Let (U, C, K) constitute an instance of MAX2SAT. Use the same widget construction as in the proof of Theorem 7. Let

$$t = 3|C| \quad (123)$$

$$\ell(\boldsymbol{\theta}) = \sum_{i=1}^t \ell(y_i - X_i; \boldsymbol{\theta}) \quad (124)$$

$$\ell^* = \min_{\boldsymbol{\theta}} \ell(\boldsymbol{\theta}) \quad (125)$$

on the constructed training set.

To prove that minimizing the bounded loss ℓ can be used to decide whether K clauses can be satisfied, first consider an intermediate clipped-loss version of the problem. In particular, let

$$\hat{\ell}(y - \hat{y}) = \frac{1}{1 - \frac{1}{2t}} \ell(y - \hat{y}) \quad (126)$$

$$\hat{\rho}(y - \hat{y}) = \min(1, \hat{\ell}(y - \hat{y})) \quad (127)$$

$$\hat{\rho}(\boldsymbol{\theta}) = \sum_{i=1}^t \hat{\rho}(y_i - X_i; \boldsymbol{\theta}) \quad (128)$$

$$\hat{\rho}^* = \min_{\boldsymbol{\theta}} \hat{\rho}(\boldsymbol{\theta}) \quad (129)$$

on the constructed training set. Note that $\hat{\ell}$ is a 1-minimal loss with $\hat{B}_1 = B_{1 - \frac{1}{2t}}$ (see Definition 4). Therefore, by Theorem 7, $\hat{\rho}^*$ can be used to decide whether K clauses can be satisfied in the original MAX2SAT instance. Recall also that $\hat{\rho}^*$ must be integer valued in this case.

Lemma 11 below shows that $\hat{\rho}^* - \frac{1}{2} \leq \ell^* \leq \hat{\rho}^*$. It then follows that for any integer n , $\hat{\rho}^* \leq n$ if and only if $\ell^* < n + \frac{1}{2}$. To see why this must hold, note: (\Leftarrow) if $\ell^* < n + \frac{1}{2}$ then $\hat{\rho}^* \leq \ell^* + \frac{1}{2} < n + 1$, hence $\hat{\rho}^* \leq n$ since $\hat{\rho}^*$ is integer valued; and also (\Rightarrow) if $\ell^* \geq n + \frac{1}{2}$ then $\hat{\rho}^* \geq \ell^* \geq n + \frac{1}{2}$, hence $\hat{\rho}^* > n$ since $\hat{\rho}^*$ is integer valued. Therefore, ℓ^* can be used to decide what integer value $\hat{\rho}^*$ achieves, which in turn can be used to decide whether K clauses can be satisfied by Theorem 7. ■

Note that bounded-loss minimization is strongly NP-hard if $B_{1 - \frac{1}{2t}}$ is polynomial in $t = 3|C|$.

Lemma 11. $\hat{\rho}^* - \frac{1}{2} \leq \ell^* \leq \hat{\rho}^*$.

Proof. First, note that for any (y, \hat{y})

$$\hat{\rho}(y - \hat{y}) - \frac{1}{2t} \leq (1 - \frac{1}{2t}) \hat{\rho}(y - \hat{y}) \quad (130)$$

$$= (1 - \frac{1}{2t}) \min(1, \hat{\ell}(y - \hat{y})) \quad (131)$$

$$= \min(1 - \frac{1}{2t}, \ell(y - \hat{y})) \quad (132)$$

$$\leq \ell(y - \hat{y}) \quad (133)$$

$$\leq \hat{\rho}(y - \hat{y}). \quad (134)$$

Therefore, for any $\boldsymbol{\theta}$

$$\hat{\rho}(\boldsymbol{\theta}) - \frac{1}{2} \leq \ell(\boldsymbol{\theta}) \leq \hat{\rho}(\boldsymbol{\theta}). \quad (135)$$

If we let $\boldsymbol{\theta}^* = \arg \min_{\boldsymbol{\theta}} \ell(\boldsymbol{\theta})$ and $\hat{\boldsymbol{\theta}} = \arg \min_{\boldsymbol{\theta}} \hat{\rho}(\boldsymbol{\theta})$, it is then immediate that

$$\hat{\rho}(\hat{\boldsymbol{\theta}}) - \frac{1}{2} \leq \hat{\rho}(\boldsymbol{\theta}^*) - \frac{1}{2} \quad (136)$$

$$\leq \ell(\boldsymbol{\theta}^*) \quad (137)$$

$$\leq \ell(\hat{\boldsymbol{\theta}}) \quad (138)$$

$$\leq \hat{\rho}(\hat{\boldsymbol{\theta}}). \quad (139)$$

■

Auxiliary References

- [26] R. Rockafellar. *Convex Analysis*. Princeton U. Press, 1970.
- [27] F. Cucker and D. Zhou. *Learning Theory: An Approximation Theory Viewpoint*. Cambridge, 2007.
- [28] Peter L. Bartlett and Shahar Mendelson. Rademacher and gaussian complexities: risk bounds and structural results. *Journal of Machine Learning Research*, 3:463–482, 2003.
- [29] K. Sturm. Generalized Orlicz spaces and Wasserstein distances for convex-concave scale functions. *Bulletin des Sciences Mathématiques*, 135:795–802, 2011.
- [30] W. Rudin. *Real and Complex Analysis*. McGraw-Hill, 3rd edition, 1987.
- [31] M. Overton and R. Womersley. Optimality conditions and duality theory for minimizing sums of the largest eigenvalues of symmetric matrices. *Mathematical Programming*, 62(2):321–357, 1993.
- [32] D. Bertsekas. *Nonlinear Programming*. Athena, 1995.
- [33] M. Primak and B. Kheifets. A modification of the inscribed ellipsoid method. *Mathematical and Computer Modelling*, 21(11):69–76, 1995.
- [34] M. Garey and D. Johnson. *Computers and Intractability: A Guide to the Theory of NP-Completeness*. W. H. Freeman and Company, 1979.
- [35] J. Håstad. Some optimal inapproximability results. *J. ACM*, 48(4):798–859, 2001.